

**Resting State Functional Connectivity
Correlates of Different
Aggression Subtypes
in Children and Adolescents with
Disruptive Behavior**

Thesis (cumulative thesis)

Presented to the Faculty of Arts and Social Sciences
of the University of Zurich
for the degree of Doctor of Philosophy

by

Julia E. Werhahn

Accepted in spring term 2019

on the recommendation of the doctoral committee:

Prof. Dr. rer. nat. Lutz Jäncke (main advisor)

Prof. Dr. Daniel Brandeis

Zurich, 2019

“When you change the way you look at things, the things you look at change.” (W. Dyer)

Contents

Summary	4
Zusammenfassung	7
1 General Introduction	11
1.1 Aggressive behavior and clinical manifestations in children and adolescents.....	11
1.2 Callous-unemotional traits and dimensions.....	12
1.3 Reactive and proactive aggression subtypes.....	14
1.4 Methods to study neural correlates of aggression subtypes – resting state fMRI.....	15
1.5 Neural alterations in aggression-related disorders.....	17
1.5.1 Neural correlates of oppositional defiant and conduct disorder.....	18
1.5.2 Neural correlates of callous-unemotional traits.....	19
1.5.3 Neural correlates of reactive and proactive aggression.....	20
1.6 Aims and hypotheses.....	21
2 Study A	23
2.1 Abstract.....	25
2.2 Introduction.....	26
2.3 Method.....	28
2.3.1 Participants.....	28
2.3.2 Clinical Assessments.....	28
2.3.3 Image Acquisition and Preprocessing.....	29
2.3.4 Regions of Interest.....	29
2.3.5 Functional Connectivity Analysis.....	30

2.4	Results.....	30
2.4.1	Sample Characteristics.....	30
2.4.2	Group Differences in Functional Connectivity.....	32
2.4.3	Functional Connectivity Correlates of Reactive versus Proactive Aggression.....	33
2.4.4	Functional Connectivity Correlates of Specific CU Dimensions.....	34
2.5	Discussion.....	36
2.6	Acknowledgments.....	39
2.7	Key points.....	40
2.8	Supplemental Information.....	41
2.8.1	Supplementary Methods and Materials.....	41
2.8.2	Supplementary Results.....	47
2.8.3	Supplemental References.....	56
3	Study B	60
3.1	Abstract.....	62
3.2	Introduction.....	63
3.3	Materials and Methods.....	65
3.3.1	Participants.....	65
3.3.2	Clinical Assessments.....	66
3.3.3	Image Acquisition.....	67
3.3.4	Preprocessing.....	67
3.3.5	Functional Connectivity Analyses.....	68
3.4	Results.....	69
3.4.1	Sample Characteristics.....	69
3.4.2	Group Differences in Functional Connectivity.....	71
3.4.3	Functional Connectivity Correlates of Reactive and Proactive Aggression.....	71

3.4.4	Functional Connectivity Correlates of Callous-Unemotional Traits.....	71
3.5	Discussion.....	73
3.5.1	Group Differences in Functional Connectivity.....	74
3.5.2	Functional Connectivity Correlates of Reactive and Proactive Aggression.....	74
3.5.3	Differential Functional Connectivity Correlates of Callous-Unemotional Traits.....	76
3.5.4	Study Strengths and Limitations.....	77
3.6	Conclusions.....	77
3.7	Acknowledgments.....	78
3.7.1	Author Disclosure Statement.....	78
3.8	Supplemental Information.....	79
3.8.1	Supplementary Materials and Methods.....	79
3.8.2	Supplementary Results.....	84
3.8.3	Supplemental References.....	91
4	General Discussion	94
4.1	Resting state fMRI in aggressive children and adolescents – main findings and future directions.....	94
4.2	Clinical implications – functional connectivity alterations during rest as a biomarker in disruptive children and adolescents?.....	103
4.3	Conclusions.....	106
	Abbreviations	108
	References	110
	Acknowledgments	134
	Curriculum Vitae	137

Summary

Clinical manifestations of severe aggressive behaviors in children and adolescents include DSM-diagnoses of conduct disorder (CD) and oppositional defiant disorder (ODD). CD characterizes frequent major violations of rules, norms, and rights. ODD manifests as pattern of angry, argumentative, and vindictive behavior. There is a considerable body of behavioral research on several aggression subtypes including reactive (RA) and proactive (PA) aggression along with callous-unemotional (CU) traits. However, knowledge of functional neural correlates at rest is limited. The majority of studies of resting state functional magnetic resonance imaging to date restricted analyses to male adolescents with CD. Additionally, RA, PA, and frequent comorbid symptoms, particularly anxiety levels, have been neglected. This dissertation aimed to fill these gaps by investigating A) seed-to-voxel cluster and B) whole-brain voxel-to-voxel resting state functional connectivity (rsFC) in 207 children and adolescents, of which 118 were cases with aggressive behavior in the clinical range including diagnoses of CD and/or ODD and 89 healthy controls, and considering RA and PA aggression subtypes along with CU traits. The present PhD project is embedded in the framework of the European multi-center FP7 projects Aggressotype and MATRICS.

In the first study, we investigated seed-based rsFC in cases compared to healthy controls, and aggression subtype-specific connectivity within cases. Recent studies reported mainly reduced rsFC in male adolescents with CD compared to healthy controls. CU and CU-related traits have been associated with predominantly increased rsFC. Affected brain regions include the posterior cingulate cortex (PCC) as part of the default mode network (DMN) and the amygdala and the anterior insula belonging to the salience network (SN). Using a common seed-based approach, regions of interest were defined in the anterior medial prefrontal cortex (amPFC) and PCC as core

regions of the DMN and in bilateral amygdala and bilateral anterior insula as parts of the SN. Compared to healthy controls, cases demonstrated reduced DMN (PCC seed) and – after controlling for ADHD symptoms – reduced SN (left anterior insula) connectivity with left hemispheric frontal clusters. Within cases, higher RA and PA scores correlated with increased connectivity in brain areas previously implicated in emotion, empathy, and cognitive control, such as the precuneus and the right caudate nucleus. All CU subdimensions correlated positively with several DMN and SN seed-based rsFC to clusters including precentral gyrus, precuneus, and cerebellar regions. These areas have been linked to emotion, empathy, moral, and self-referential processes. This first study thus revealed aggression subtype-specific rsFC patterns, extending the knowledge of neural networks involving two core regions implicated in aggression-related disorders to distinct forms of disruptive behavior in children and adolescents.

In the second study, we extended previous efforts by investigating unrestricted graph theory based whole-brain voxel-to-voxel rsFC. By calculating the Intrinsic Connectivity Contrast (ICC) as global connectivity strength of each voxel with all other voxels, we measured changes in network centrality. Additionally, analysis of Integrated Local Correlation (ILC) examined changes in local coherence by integrating voxel-wise spatial correlations. Besides ADHD symptoms, co-occurring anxiety levels were considered. Cases showed altered rsFC including the medial frontal cortex compared to controls, but only when anxiety and not ADHD symptoms were controlled for. Importantly, most of the aggression subtype-specific voxel-wise rsFC patterns within cases depended on the additional control of both ADHD and anxiety symptoms. Both RA and PA symptoms related to connectivity alterations including parietal areas and distinct connectivity patterns in central gyrus and precuneus, which seem to be involved in aggression, emotion, and cognitive control-related processes. CU traits correlated with patterns including temporal and cerebellar areas implicated in empathy, emotion, and reward-related activity. This hypothesis-free rsFC study further supported an aggression subtype-specific approach and underlined the crucial role of considering comorbid symptoms in such analyses.

In summary, this dissertation project points to the sensitivity of rsFC to developmental forms of aggression, and to the importance of considering different manifestations of aggression in children and adolescents with disruptive behavior. Firstly, both seed- and unrestricted voxel-based whole-brain analysis approaches showed alterations in rsFC including frontal brain areas in cases compared to healthy controls, which partly depended on the additional control of either ADHD or anxiety symptoms. Secondly, both analysis approaches revealed altered and predominantly distinct rsFC related to RA and PA subtypes along with CU traits within cases. Importantly, comorbid ADHD and anxiety symptoms were crucial in identifying aggression subtype-specific voxel-wise rsFC alterations. Taken together, the significant aggression subtype-specific seed-based and voxel-wise rsFC results in a large sample of children and adolescents with aggression elucidated that distinct though overlapping brain connectivity measures can disentangle distinct forms of aggressive behaviors and underline a dimensional rather than categorical approach. Our results could have clinical implications, as they might provide support for personalized diagnostic and treatment of aggression-related disorders tailored to the specific neural manifestation of aggressive behavior.

Zusammenfassung

Schwerwiegend aggressives Verhalten von Kindern und Jugendlichen kann anhand von DSM-Diagnosen wie der Störung des Sozialverhaltens (CD) und der Oppositionellen Störung des Sozialverhaltens (ODD) klinisch erfasst werden. CD beschreibt wiederholte erhebliche Verletzung von Regeln, Normen und Rechten. ODD zeigt sich anhand von wütenden, streitsuchenden und rachsüchtigen Verhaltensmustern. Von zentraler Bedeutung sind dabei einige Subtypen von Aggression wie reaktive (RA) und proaktive (PA) Aggression sowie verschiedene Ausprägungen eingeschränkter prosozialer Emotionalität (CU-Traits). Während es auf Verhaltensebene immer mehr Forschung zu diesen Aggressionsformen gibt, ist das Wissen über funktionelle neuronale Korrelate während des Ruhezustandes noch begrenzt. Bis heute beschränkten Studien über den Ruhezustand des Gehirns, gemessen mit funktioneller Magnetresonanztomographie, ihre Analysen meistens auf männliche Jugendliche mit CD. Ausserdem blieben RA, PA sowie komorbide Symptome, vor allem Ausprägungen von Angst, bislang unbeachtet. Diese Dissertation setzte sich das Ziel, ebenjene Forschungslücken zu schliessen. Wir untersuchten demzufolge A) seedbasierte funktionelle Konnektivitäten im Ruhezustand (rsFC) und B) rsFC einzelner Voxel mit allen übrigen Voxeln im Gehirn von 207 Kindern und Jugendlichen. Dabei zeigten 118 Studienteilnehmende ausgeprägtes aggressives Verhalten mit Werten im klinischen Bereich und hatten zumeist auch Diagnosen von CD und/oder ODD erhalten. 89 waren gesunde Kontrollprobanden. Wir berücksichtigten RA und PA Aggressionssubtypen nebst CU-Traits. Das vorliegende Dissertationsprojekt ist in die europäischen Multicenter FP7-Projekte Aggressotype und MATRICS eingebettet.

In der ersten Studie wurden Unterschiede zwischen aggressiven und gesunden Kindern und Jugendlichen hinsichtlich seedbasierter rsFC

untersucht, und ob sich diese abhängig von der Aggressionsform innerhalb der aggressiven Gruppe unterscheidet. Bisherige Studien berichteten vorwiegend reduzierte rsFC bei männlichen Jugendlichen mit CD im Vergleich zu gesunden. CU- und CU-ähnliche Traits wurden mit meist erhöhter rsFC in Verbindung gebracht. Zu den betroffenen Gehirnarealen zählen der posteriore cinguläre Cortex (PCC), der zum Default-Mode-Netzwerk (DMN) gehört, sowie Amygdala und anteriore Insel, Bestandteile des Salienz-Netzwerks (SN). Mittels der verbreiteten seedbasierten Herangehensweise wurden Bereiche von Interesse im anterioren medialen Präfrontalkortex (amPFC) und im PCC (Kernregionen des DMN), sowie in bilateraler Amygdala und bilateraler anteriorer Insel (Bestandteile des SN) erstellt. Verglichen mit den gesunden zeigten die aggressiven Kinder und Jugendliche eine verringerte Konnektivität mit linkshemisphärischen frontalen Clustern, zum einen mit Bereich von Interesse im DMN (PCC seed), zum anderen nach Kontrolle für ADHS-Symptome im SN (linksseitige anteriore Insel). Innerhalb der aggressiven Studienteilnehmenden korrelierten höhere RA- und PA-Ausprägungen mit stärkeren Konnektivitäten in Gehirnarealen wie etwa dem Precuneus und dem rechten Nucleus Caudatus, die vormalig bei Emotion, Empathie und kognitiver Kontrolle involviert waren. Alle CU-Subdimensionen korrelierten positiv mit einigen DMN und SN seedbasierten rsFC mit Clustern, die den Gyrus präcentralis, Precuneus und zerebellare Regionen beinhalten. Diese Areale wurden mit emotionalen, empathischen, moralischen, und selbstreferenziellen Prozessen in Verbindung gebracht. Diese erste Studie zeigte somit aggressionssubtypspezifische rsFC Muster, die das Wissen über die bei Aggressionsstörungen beteiligten neuronalen Netzwerke, welche zwei Kernregionen miteinschließt, auf spezifische Formen aggressiven Verhaltens von Kindern und Jugendlichen ausweitete.

In der zweiten Studie erweiterten wir vorherige Bestrebungen, indem wir uneingeschränkte, graphentheoretische rsFC zwischen einzelnen Voxeln im gesamten Gehirn untersuchten. Durch die Berechnung des «Intrinsic-Connectivity-Contrast» (ICC), der globalen Konnektivitätsstärke eines jeden Voxels mit allen anderen Voxeln, erfassten wir Veränderungen in der Netzwerkzentralität. Zusätzlich untersuchten wir mit der «Integrated-

Local-Correlation» (ILC) Veränderungen in lokaler Kohärenz über die Integration voxelbasierter räumlicher Korrelationen. Neben ADHD-Symptomen wurden auch komorbide Angstausprägungen berücksichtigt. Aggressive Kinder und Jugendliche zeigten nur dann abweichende rsFC im Vergleich zu gesunden Teilnehmenden, etwa im medialen Frontalkortex, wenn für Angst- aber nicht für ADHD-Symptome kontrolliert wurde. Von zentraler Bedeutung ist, dass die meisten aggressionssubtypspezifischen rsFC-Resultate innerhalb der aggressiven Studienteilnehmenden von der zusätzlichen Kontrolle für sowohl ADHD- also auch Angstsymptome abhängen. RA- und PA-Symptome standen beide in Zusammenhang mit abweichenden Konnektivitäten in parietalen Arealen und mit unterschiedlichen Konnektivitätsmustern im zentralen Gyrus und Precuneus. Diese Regionen scheinen bei aggressionsbezogenen, emotionalen und kognitiven Kontrollprozessen beteiligt zu sein. CU-Traits korrelierten mit Konnektivitätsmustern in temporalen und zerebralen Gehirnarealen, die mit Empathie, Emotion, und Belohnung in Verbindung gebracht werden. Diese hypothesenfreie rsFC-Studie untermauerte ebenfalls eine entsprechend den Aggressionssubtypen ausgerichtete Vorgehensweise, und betonte die zentrale Rolle des Einbezugs komorbider Symptome in ebensolchen Analysen.

Zusammenfassend weist dieses Dissertationsprojekt auf die Sensitivität von rsFC für entwicklungsbedingte Aggressionsformen hin und betont die Bedeutsamkeit, unterschiedliche Manifestationen von Aggression bei Kindern und Jugendlichen mit Störungen des Sozialverhaltens oder ausgeprägter Aggression zu berücksichtigen. Zunächst zeigten sowohl seed- als auch unbeschränkte voxelbasierte Analyseformen abweichende rsFC von aggressiven im Vergleich zu gesunden Kindern und Jugendlichen. Diese beinhaltete frontale Gehirnbereiche und hing teilweise von der zusätzlichen Kontrolle für entweder ADHD- oder Angstsymptome ab. Zum anderen zeigten beide analytischen Herangehensweisen abweichende und überwiegend spezifische rsFC in Zusammenhang mit den unterschiedlichen RA- und PA-Subtypen sowie CU-Traits innerhalb der aggressiven Studienteilnehmenden. Von zentraler Bedeutung ist die wesentliche Rolle, die komorbiden ADHD- und Angstsymptomen bei der Identifikation

aggressionssubtypspezifischer Auffälligkeiten in voxelbasierter rsFC zukam. Insgesamt zeigten diese signifikanten, sich abhängig von den Aggressionssubtypen unterscheidenden seed- und voxelbasierten Konnektivitätsmuster in einer grossen Stichprobe von aggressiven Kindern und Jugendlichen, dass verschiedene und dabei doch auch überlappende Messmethoden von Gehirnkonnektivitäten die spezifischen Formen aggressiven Verhaltens entflechten können. Darüber hinaus unterstützen diese Resultate eine dimensionale statt kategoriale Herangehensweise. Unsere Ergebnisse könnten insofern klinische Implikationen haben, als dass sie eine entsprechend den unterschiedlichen neuronalen Manifestationen von aggressivem Verhalten zugeschnittene, personalisierte Diagnostik und Behandlung von Aggressionsstörungen unterstützen.

1 General Introduction

1.1 Aggressive behavior and clinical manifestations in children and adolescents

From an evolutionary psychological perspective, there are different hypotheses on the origin of aggression. Aggression not only reflects problematic behavior but often an adaptive and context-dependent way to deal with challenging situations (Buss & Shackelford, 1997). Aggression-related behavior can be understood as an attempt to deal with perceived threat or as resulting from impaired learning to inhibit instinctive fight/flight responses (Fonagy & Luyten, 2017). Physical aggression was shown to start within the first year after birth. Thus, longitudinal studies started with birth cohorts and found an increasing frequency of physical aggression until the age of three to four years, which usually declines with physical growth (Tremblay, Vitaro, & Côté, 2018). However, aggression levels of most physically aggressive children show a small increase during adolescence, followed by a decrease during adulthood. Stimulation-seeking (Zuckerman & Neeb, 1979) might be an attempt of antisocial individuals to increase low autonomic arousal levels (van Goozen, Fairchild, Snoek, & Harold, 2007). From a longitudinal perspective, high levels of stimulation-seeking and fearlessness at the age of three years have been related to aggression levels eight years later (Raine, Reynolds, Venables, Mednick, & Farrington, 1998).

Aggression-related disorders in childhood and adolescence are associated with poor outcomes including criminal behavior, substance use problems, and antisocial personality disorders (Erskine et al., 2016) besides a substantial contribution to the global burden of disease (Erskine et al., 2014). There is ongoing research on developmental pathways to aggression-related disorders involving discussions on the impact of the age

of onset, for instance (Frick, 2012; Puzzo, Smaragdi, Gonzalez, Martin-Key, & Fairchild, 2016). Recent research suggests an important interplay of genetic predispositions like specific candidate genes and environmental factors including parental monitoring (Salvatore & Dick, 2016). From a clinical perspective, recurrent angry, arguing, and revengeful mood and behavior, leading to individual distress in others or negatively effect daily interactions or functioning in social environments, can manifest itself as a diagnosis as oppositional defiant disorder (ODD) in the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) (American Psychiatric Association, 2013). If three or more criteria including severe violation of norms, rules, and rights within the past year are present, a diagnosis of conduct disorder (CD) is given. ODD and CD are among the most common psychiatric disorders in childhood and adolescence with a world-wide prevalence of 5.7% (3.6% for ODD and 2.1% for CD) (Polanczyk, Salum, Sugaya, Caye, & Rohde, 2015). According to the review by Waschbusch (Waschbusch, 2002), 36% of boys and 57% of girls with conduct problems have comorbid hyperactive-impulsive-inattention symptoms. Particularly the combination of an early onset of CD problems in childhood and impaired self-control is prone to a longer persistence of problematic behaviors with a peak of antisocial behavior in adolescence (Moffitt, 1993). Moreover, anxiety symptoms are frequently comorbid in children with conduct problems (Frick, 2012; Frick, Ray, Thornton, & Kahn, 2013). The effectiveness of treating aggression-related disorders in children and adolescents is small (Bakker, Greven, Buitelaar, & Glennon, 2017). As delineated in the following, a growing body of evidence elucidated the importance to differentiate between different forms of disruptive behavior.

1.2 Callous-unemotional traits and dimensions

The role of callous-unemotional (CU) traits, newly added as a specifier for CD in the DSM-5 (American Psychiatric Association, 2013), has been frequently investigated (Frick, 2016). CU traits are also named “with limited prosocial emotions” to reduce stigmatizing (Blair, Leibenluft, & Pine, 2014). Recent reviews suggested high levels of heritability (Raine, 2018) and that

CU traits might predict future antisocial behavior (Waller & Hyde, 2017), even during childhood (Dadds, Fraser, Frost, & Hawes, 2005), and adult psychopathy (Frick & White, 2008). Childhood CU traits and adult psychopathic features seem related (Frick et al., 2013; Frick & White, 2008). The interaction of inherited emotion-related deficits and fearlessness with parental practices might contribute to an increasing risk for early CU traits (Waller & Hyde, 2018). CU traits have been associated with fearlessness and higher unconcern about punishment (Fanti, Panayiotou, Lazarou, Michael, & Georgiou, 2016), impaired emotion recognition skills, particularly for fear (Muñoz, 2009; Sylvers, Brennan, & Lilienfeld, 2011), and reward-oriented behavior suggested to lead to a higher risk for most severe, frequent, and instrumental violent behavior (Reidy et al., 2017). Higher levels of CU traits related to more severe conduct problems in childhood (Longman, Hawes, & Kohlhoff, 2016) and seem to predict antisocial behavior (Dadds et al., 2005), delinquent behavior, and a diagnosis of antisocial behavior disorder in adulthood (McMahon, Witkiewitz, & Kotler, 2010). Despite stability during childhood peculiarly in fearless children (Goffin, Boldt, Kim, & Kochanska, 2017) and during adolescence (Eisenbarth, Demetriou, Kyranides, & Fanti, 2016; Frick et al., 2013; Reidy et al., 2017), CU traits tend to decrease from childhood to adolescence. As such, a minor decrease of CU traits during childhood correlated with an increased risk for externalizing problems in adolescence (Muratori et al., 2016). A recent review (Moul, Hawes, & Dadds, 2016) shed light on neurobiological alterations in children with conduct problems and CU traits. For instance, reduced functioning of the oxytocin system and serotonin neurotransmission seems related to poor emotion-recognition, which might in turn contribute to an impaired development of empathy.

In recent years, a growing body of research pointed to differentiating between dimensions of CU traits (Kimonis et al., 2008; Latzman, Lilienfeld, Latzman, & Clark, 2013; Pechorro, Ray, Gonçalves, & Jesus, 2017; Pihet, Etter, Schmid, & Kimonis, 2015) and CU-related traits (Salihovic & Stattin, 2017). Youths with CU and CU-related traits show differing anxiety levels that seem related to differential behaviors (Fanti, Demetriou, & Kimonis, 2013; Kimonis, Frick, Cauffman, Goldweber, & Skeem, 2012) like impaired

emotion recognition skills (Dadds, Kimonis, Schollar-Root, Moul, & Hawes, 2017) and higher levels of aggression-related behaviors (Fanti et al., 2013; Guelker, Barry, Barry, & Malkin, 2014).

1.3 Reactive and proactive aggression subtypes

Reactive (RA) and proactive (PA) aggression symptoms (Dodge & Coie, 1987; Vitiello & Stoff, 1997) also characterize important aspects of aggressive behavior. RA symptomatology describes aggressive reactions to internal or external provocation (Smeets et al., 2016), while PA specifies instrumental, planned aggressive behavior in order to influence others (Dodge & Coie, 1987). There is evidence for a differing heritability of RA and PA symptomatology (Waltes, Chiocchetti, & Freitag, 2016). For instance, harsh parenting has predicted both aggression forms, while negative emotionality was shown to predict RA but not PA behavior in children (Vitaro, Barker, Boivin, Brendgen, & Tremblay, 2006). High levels of RA and PA symptoms seem to increase in early and decline in late adolescence (Tremblay et al., 2018).

On the one hand, existing behavioral literature suggests a substantial correlation of the proposed RA and PA subtypes (Card & Little, 2006; Polman, Orobio De Castro, Koops, Van Boxtel, & Merk, 2007; Poulin & Boivin, 2000) and commonalities (Cima, Raine, Meesters, & Popma, 2013; Marsee et al., 2011). For instance, both have been associated with CU traits (Feilhauer, Cima, & Arntz, 2012; Kimonis et al., 2008; Pechorro et al., 2017) and CU-related traits (Blais, Solodukhin, & Forth, 2014; Perenc & Radochonski, 2014) and with increased CD symptoms and CU traits in adolescents (Eisenbarth et al., 2016). On the other hand, there is support for their differentiation on a psychophysiological (Scarpa, Haden, & Tanaka, 2010) and behavioral level. While RA behavior has been linked to impulsivity and social anxiety, PA symptoms seem related to delinquency and hyperactivity (Raine, Dodge, Loeber, Gatzke-Kopp, Lynam, Stouthamer-Loeber, et al., 2006). High levels of RA symptoms seem to increase the risk for higher internalizing problems (Fite, Rubens, Preddey, Raine, & Pardini, 2014) and related to impulsivity (Connor, Steingard, Anderson, & Melloni,

2003; Fite, Stoppelbein, & Greening, 2009; Urben et al., 2018) and suicidal behaviors (Hartley, Pettit, & Castellanos, 2016), and are thought to positively predict CU-related traits in early adolescence (Bamvita et al., 2017). In contrast, PA symptoms seem associated with reduced moral-related processes in early childhood (Jambon & Smetana, 2018), conduct problems (Smeets et al., 2016), and CU or CU-related traits (Fite et al., 2009; Marsee et al., 2011; Marsee & Frick, 2007; Urben et al., 2018; White, Gordon, & Guerra, 2015). Psychophysiological research in children (Scarpa et al., 2010) and adolescents (Gao, Tuvblad, Schell, Baker, & Raine, 2015) reported both distinct and inconsistent findings (Fanti, 2016) regarding the RA/PA-related differentiation.

Taken together, considering RA and PA behaviors seems inevitable when subtyping aggression-related disorders in children and adolescents.

1.4 Methods to study neural correlates of aggression subtypes – resting state fMRI

During magnetic resonance imaging (MRI), changes in physiological events are reflected by changes in blood-oxygen-level-dependent (BOLD) signals (Ogawa, Lee, Kay, & Tank, 1990). Studies of functional MRI (fMRI) assume that active brain regions are characterized by a local change in the amount of measured deoxygenated hemoglobin in the blood (Huettel, Song, & McCarthy, 2009). Neural activity is measured indirectly and non-invasively with good spatial resolution as the difference between oxygenated and deoxygenated blood by recording changes in the MR signal emerging as radiofrequency waves. The process of excitation is described as the application of a radio frequency pulse to protons spinning parallel to a strong magnetic field (longitudinal magnetization), which leads to high-energy states in proton spins (transverse magnetization). Stopping the pulse leads to a process of relaxation, as the protons return to a low-energy state. The time of recovering the longitudinal magnetization is called longitudinal relaxation (T1). Afterwards, the signal of transverse components decays, measured as transverse relaxation time (T2). The BOLD-contrast during fMRI is based on the time constant T_2^* , which

delineates the decay of transverse magnetization due to interactions of rotating spins precessing at higher or lower frequencies and spatial inhomogeneities in the local magnetic field (Huettel et al., 2009).

Biswal and colleagues (Biswal, Yetkin, Haughton, & Hyde, 1995) were the first describing resting state fMRI (rs-fMRI) when they detected that low-frequency time courses of brain regions during hand movement are also detectable during periods of rest, along with further regions relating to motor function. There is evidence for distinct highly functionally connected brain regions active in the absence of goal-directed behaviors, known as resting state networks (RSNs) (Damoiseaux et al., 2006; Fox, Spreng, Ellamil, Andrews-Hanna, & Christoff, 2015; Greicius, Krasnow, Reiss, & Menon, 2003; Raichle et al., 2001). For instance, the default mode network (DMN) consists of a distinct set of functionally highly connected brain regions activated during rest (Andrews-Hanna, Reidler, Sepulcre, Poulin, & Buckner, 2010; Fox et al., 2015; Raichle et al., 2001) across all age groups (Mak et al., 2017) including children (Thomason et al., 2011). The connectivity strength of RSNs was shown to exhibit moderate to high test-retest reliability across subjects and runs with varying acquisition parameters (van Dijk et al., 2010). An fMRI acquisition method emerging in recent years collects multiple instead of single echo images for each slice. This allows for modelling the T_2^* signal decay for each voxel, which in turn enables to differentiate between BOLD and artifact signal (Kundu et al., 2017). For instance, the effect of motion-related noise on DMN rsFC could be minimized by using a multi-echo compared to a single-echo approach (Dipasquale et al., 2017).

There are different ways to analyze rs-fMRI data. The hypothesis-driven seed-based approach is among the most frequently applied methods for analyzing rsFC (Biswal et al., 1995; Fox et al., 2005; Greicius et al., 2003; Whitfield-Gabrieli & Ford, 2012). After a priori definition of regions of interest (ROIs), the correlation between BOLD time courses from a seed and whole-brain voxel clusters is calculated at a predefined threshold (Lee, Smyser, & Shimony, 2013). In order to further explore rs-fMRI data, unrestricted, hypothesis-free, and graph theory-based approaches include the Intrinsic Connectivity Contrast (ICC) and the Integrated Local

Correlation (ILC) (Whitfield-Gabrieli & Nieto-Castanon, 2012). ICC enables to analyze rsFC alterations in the whole brain without an a priori definition of ROIs (Martuzzi et al., 2011). Using ICC, the global correlation strength of voxel-to-voxel connectivity patterns is calculated, which measures changes in network centrality (Whitfield-Gabrieli & Nieto-Castanon, 2012). Regarding local voxel-wise rsFC patterns, ILC enables to examine changes in local coherence by integrating spatial correlations of each voxel without the need for a specified finite neighborhood for the integration, as in other local coherence approaches (Deshpande, LaConte, Peltier, & Hu, 2009).

Numerous studies provide evidence for psychiatric disorders to be associated with altered rsFC (Woodward & Cascio, 2015). Thus, studies of rs-fMRI may attribute to diagnosis and personalized treatment by exploring relations to psychiatric disorders (Barch, 2017; Whitfield-Gabrieli & Ford, 2012).

1.5 Neural alterations in aggression-related disorders

There is evidence for neural alterations on both a structural and functional level, with functional ones at rest and in tasks probing aggression-related processing. Antisocial and disruptive youths have been reported to show less punishment and higher immediate reward oriented behavior and thus deficit functioning of associated brain areas (Byrd, Loeber, & Pardini, 2014) including (pre-) frontal regions like the medial prefrontal cortex along with temporal alterations (Alegria, Radua, & Rubia, 2016). Another review and meta-analysis documented both structural and functional neural abnormalities including the left insula (Raschle, Menks, Fehlbaum, Tshomba, & Stadler, 2015). Interestingly, ADHD and disruptive behavior disorders seem to exhibit overlapping neural impairments in cognitive control networks which include the orbitofrontal cortex (OFC), cingulate, and insular cortices (Puiu et al., 2018). Given the heterogeneity within aggression-related disorders associated with different neural alterations, it is of particular concern to differentiate between the distinct manifestations and traits also on a neural level (Blair, Veroude, & Buitelaar, 2016).

1.5.1 Neural correlates of oppositional defiant and conduct disorder

In CD and ODD, disrupted functioning of frontal, striatal, and limbic regions has been associated with altered emotional, reward, and punishment processing and related regulatory mechanisms (Blair et al., 2016; Noordermeer, Luman, & Oosterlaan, 2016). For instance, besides alterations in frontal brain regions including the OFC implicated in disrupted reward-related behavior also in ADHD (Von Rhein et al., 2015), functioning of the amygdala (Holz et al., 2016) and anterior insular cortex correlated with different forms of empathic processes, which are impaired in adolescents with conduct problems (Blair et al., 2016). As reviewed recently, abnormal fear conditioning and, closely linked, punishment processing in ODD and CD have been associated with altered amygdala activity, which in turn seems implicated in reward-related processes (Matthys, Vanderschuren, & Schutter, 2013). Moreover, the impaired cognitive regulation of emotional behaviors is thought to be associated with structural and functional alterations in paralimbic regions including the OFC and cingulate cortices. A recent meta-analysis identified reduced gray matter volumes in brain regions such as left amygdala, right insula (Rogers & De Brito, 2016), and left central gyrus along with the posterior cingulate cortex (PCC) depending on the CD severity (Smaragdi et al., 2017) in youths.

During rest, many of the networks activated or deactivated in tasks can also be observed using fMRI. Recent studies in male adolescents with CD suggested mainly diminished resting state functional connectivity (rsFC) (Aghajani et al., 2017; Broulidakis et al., 2016; Lu, Zhou, Wang, Xiang, & Yuan, 2017; Lu et al., 2015; Zhou et al., 2016) and activity (Lu et al., 2015; Wu, Zhang, Dong, Wang, & Yao, 2017). For instance, male adolescents with CD demonstrated reduced rsFC between the core regions of the DMN, anterior medial prefrontal cortex (amPFC) and PCC, compared to healthy control subjects only after controlling for ADHD symptoms, as ADHD symptoms correlated positively with rsFC within the DMN (Broulidakis et al., 2016). Further studies found increased (Pu et al., 2017) but mainly reduced rsFC including areas of the DMN (Lu, Zhou, Wang, et al., 2017; Lu et al., 2015; Zhou et al., 2016) and the somatosensory network (Lu et al., 2015).

Diminished neural activity during rest in brain regions has been reported in parts of the salience network (SN) (Zhou, Yao, Fairchild, Zhang, & Wang, 2015) along with cerebellar and parietal areas (Wu et al., 2017). Another recent study in male and female patients with CD documented altered spontaneous activity including a reduced amplitude of low frequency fluctuations in left inferior/middle temporal gyrus (ITG/MTG) in males compared to females (Cao, Sun, Dong, Yao, & Huang, 2018). Impulsivity scores have been related to mainly increased rsFC in SN (Pu et al., 2017) and DMN regions (Lu, Zhou, Wang, et al., 2017; Lu, Zhou, Zhang, Wang, & Yuan, 2017; Shannon et al., 2011) also found in boys with CD and ADHD (Uytun et al., 2016).

In sum, most of rs-fMRI studies to date exclusively included male adolescents with CD, had sample sizes for cases of $N < 60$, differing methodological approaches, and disregarded distinct heterogeneous manifestations of aggression-related disorders such as RA and PA symptomatology. This makes it difficult to draw reliable conclusions on rs-fMRI-derived neural alterations children and adolescents.

1.5.2 Neural correlates of callous-unemotional traits

Youths presenting with CU traits have been characterized by reduced functioning of brain regions related to reinforcement and emotional empathy, such as amygdala (Blair et al., 2014; Herpers, Scheepers, Bons, Buitelaar, & Rommelse, 2014) and anterior insula (Michalska, Zeffiro, & Decety, 2016; Reidy et al., 2017). A recent study reported that CU traits predicted a reduced cingular-limbic functional connectivity when viewing harm to others (Yoder, Lahey, & Decety, 2016), while in another study CU traits correlated with reduced cingular response to fearful faces (Szabó et al., 2017). Psychopathic traits in adulthood have been associated with (para-)limbic abnormalities (Kiehl, 2006) and the amygdala has been suggested as core region (Blair, 2017). However, there is growing evidence for neural alterations beyond (para-)limbic regions (Griffiths & Jalava, 2017). On a structural level, CU traits related to alterations in temporal regions (Cohn et al., 2016; Fairchild et al., 2015; Wallace et al., 2014) along

with frontal (Fairchild et al., 2013; Umbach, Berryessa, & Raine, 2015) and parietal areas (Yang et al., 2015).

Only a limited number of rsFC studies considered CU traits in children and adolescents, thereby mostly including male adolescents with CD. There are reports of CU and CU-related dimension-specific increases in regions of the DMN (Cohn et al., 2015; Thijssen & Kiehl, 2017) along with the SN and other resting state networks (RSNs) (Thijssen & Kiehl, 2017). Juvenile offenders with CD and differing levels of CU(-related) traits were shown to exhibit differing rsFC patterns including clusters in fronto-limbic and fronto-parietal areas (Aghajani et al., 2016, 2017). Likewise, there is evidence for psychopathic traits in adults to be considered. For instance, incarcerated male adults exhibited altered connectivity in multiple RSNs related to the interpersonal/affective factor of psychopathy (Espinoza et al., 2018). Another study reported an increase and decrease of rsFC depending on the psychopathic trait in brain regions including anterior insula and PCC (Philippi et al., 2015). Despite behavioral evidence for the crucial, moderating role of co-occurring anxiety levels on a behavioral level (Fanti et al., 2013), only one rsFC study considered these and documented an interaction effect of anxiety and psychopathy levels in adults (Motzkin, Newman, Kiehl, & Koenigs, 2011).

1.5.3 Neural correlates of reactive and proactive aggression

To date, few magnetic resonance imaging (MRI) studies have taken RA and PA manifestations of aggression into account. RA symptomatology might result from failed inhibition of aggressive impulses by executive brain systems (Blair, 2001). Blair (Blair, 2016) suggested frontal areas and the anterior insula cortices involved, which in turn might be implicated in reinforcement-based decision-making. Furthermore, RA behavior is thought to be mediated by the so-called acute threat response system. This neural circuit includes amygdala, hypothalamus, and periaqueductal gray, and seems modulated by prefrontal areas like the OFC (Blair, 2004; Blair, 2010). According to a recent review of fMRI studies on RA symptomatology, pathological aggression has been suggested to be associated with

hyperresponsivity in emotion-related neural circuits including the amygdala and insula and hyporesponsivity in an emotion regulation circuitry that comprises cingular and parietal areas like the precuneus (Fanning, Keedy, Berman, Lee, & Coccaro, 2017). Children with CD symptoms were shown to exhibit a positive correlation between posterior insula response to emotional empathy stimuli and RA scores (Michalska et al., 2016). With respect to PA symptomatology, amygdala dysfunction might contribute to impaired development of moral-related behaviors (Blair, 2004). Yet, there is evidence that both RA and PA symptoms might relate to altered amygdala functioning. For instance, a reduced amygdala response to fear in children with high CU traits predicted PA levels (Lozier, Cardinale, VanMeter, & Marsh, 2014). During rs-fMRI, male adult offenders exhibited a positive correlation between RA and PA scores and increased left amygdala rsFC with medial prefrontal cortex, and a RA symptoms-related decrease of left amygdala rsFC with right posterior insula (Siep et al., 2018). Furthermore, higher levels of PA and CU-related traits in adolescence have been associated with lower right amygdala volume in adulthood (Pardini, Raine, Erickson, & Loeber, 2014).

To the best of our knowledge, no rsFC study in aggressive children and adolescents examined the role of RA and PA subtypes to date.

1.6 Aims and hypotheses

This dissertation project aimed to extend the knowledge of aggression-related neural networks to distinct subtypes of aggression using rs-fMRI data derived from children and adolescents in a multi-center study. To date, studies investigating rsFC in aggressive youth restricted analysis predominantly to male adolescents with pure CD. Hence, little is known about further and specific manifestations of aggression. Particularly RA and PA subtypes along with comorbid ADHD and anxiety symptoms have been widely neglected. The current thesis attempted to fill these gaps in literature. We thus investigated rsFC in male and female children and adolescents with aggressive behavior in the clinical range including diagnoses of CD/ODD compared to controls thereby considering comorbid

symptoms. Most notably, we aimed to elucidate connectivity alterations related to RA and PA symptoms along with CU traits within cases.

In a first study, we investigated seed-based rsFC with whole-brain voxel clusters using literature-based a priori defined ROIs within the DMN and the SN. We expected reduced seed-based connectivity in cases compared to healthy controls. We considered comorbid ADHD symptoms and assumed an increase in rsFC with higher ADHD scores. Furthermore, we hypothesized that CU subdimensions would be associated with distinct and increased rsFC within cases. For RA and PA symptomatology, we tentatively speculated to find differing rsFC patterns particularly for amygdala and anterior insula seeds.

In a second study using the same sample and data, we extended previous efforts by using hypothesis-free voxel-to-voxel whole-brain approaches. ICC was employed to assess changes in global correlation strength and ILC to calculate alterations in local coherence. For all analyses, we took into account both comorbid ADHD and anxiety symptoms and expected them to partly modulate (enhance or suppress) aggression-related connectivity patterns. We hypothesized that voxel-wise rsFC alterations are found in cases compared to controls, and that these include a reduced recruitment of frontal areas. Within cases, we expected to find patterns including the precuneus related to RA and PA symptoms along with further temporal, parietal, and frontal regions associated with CU traits beyond (para-)limbic regions.

2 Study A

AGGRESSION SUBTYPES RELATE TO DISTINCT RESTING STATE FUNCTIONAL CONNECTIVITY IN DISRUPTIVE CHILDREN AND ADOLESCENTS

Julia E Werhahn^{1,2}, Susanna Mohl¹, David Willinger^{1,2}, Lukasz Smigielski^{1,2}, Alexander Roth¹, Christoph Hofstetter¹, Philipp Stämpfli³, Isabelle Häberling¹, Jilly Naaijen^{4,5}, Leandra M Mulder^{4,5}, Jeffrey C Glennon⁴, Pieter J Hoekstra⁶, Andrea Dietrich⁶, Renee Kleine Deters⁶, Pascal M Aggensteiner⁷, Nathalie E Holz⁷, Sarah Baumeister⁷, Tobias Banaschewski⁷, Melanie C Saam⁸, Ulrike M E Schulze⁸, David J Lythgoe⁹, Arjun Sethi⁹, Michael Craig⁹, Mathilde Mastroianni¹⁰, Ilyas Sagar-Ouriaghli¹⁰, Paramala J Santosh¹⁰, Mireia Rosa¹¹, Nuria Bargallo¹², Josefina Castro-Fornieles¹³, Celso Arango¹⁴, Maria J Penzol¹⁴, Marcel P Zwiers⁵, Barbara Franke^{15,16}, Jan K Buitelaar^{4,17}, Susanne Walitza^{1,2,18}, Daniel Brandeis^{1,2,7, 18}

¹Department of Child and Adolescent Psychiatry and Psychotherapy, University Hospital of Psychiatry Zurich, University of Zurich, Zurich, Switzerland

²Neuroscience Center Zurich, University and ETH Zurich, Zurich, Switzerland

³MR-Center of the Department of Psychiatry, Psychotherapie and Psychosomatics and the Department of Child and Adolescent Psychiatry, Psychiatric Hospital, University of Zurich, Zurich, Switzerland

⁴Radboud University Medical Center, Donders Institute for Brain, Cognition and Behavior, Department of Cognitive Neuroscience, Nijmegen, The Netherlands

⁵Radboud University, Donders Institute for Brain, Cognition and Behavior, Centre for Cognitive Neuroimaging, Nijmegen, The Netherlands

⁶University of Groningen, University Medical Center Groningen, Department of Child and Adolescent Psychiatry, Groningen, The Netherlands

⁷Department of Child and Adolescent Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim/Heidelberg University, Mannheim, Germany

⁸Department of Child and Adolescent Psychiatry/Psychotherapy, University Hospital, University of Ulm, Germany

⁹Department of Neuroimaging, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

¹⁰Department of Child Psychiatry, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

¹¹Child and Adolescent Psychiatry Department, Hospital Clinic of Barcelona, IDIBAPS, Barcelona, Spain

¹²Clinic Image Diagnostic Center (CDIC), Hospital Clinic of Barcelona; Magnetic Resonance Image Core Facility, IDIBAPS, Barcelona, Spain

¹³Child and Adolescent Psychiatry and Psychology Department, 2017SGR881, Institute Clinic of Neurosciences, Hospital Clinic of Barcelona, CIBERSAM, IDIBAPS, Department of Medicine, University of Barcelona, Barcelona, Spain

¹⁴Child and Adolescent Psychiatry Department, Hospital General Universitario Gregorio Marañón School of Medicine, Universidad Complutense, IiSGM, CIBERSAM, Madrid, Spain

¹⁵Radboud University Medical Center, Donders Institute for Brain, Cognition and Behavior, Department of Human Genetics, Nijmegen, The Netherlands

¹⁶Radboud University Medical Center, Donders Institute for Brain, Cognition and Behavior, Department of Psychiatry, Nijmegen, The Netherlands

¹⁷Karakter Child and Adolescent Psychiatry University Center, Nijmegen, The Netherlands

¹⁸Zurich Center for Integrative Human Physiology, University of Zurich, Zurich, Switzerland

Running title: Aggression subtype-specific connectivity

Keywords: reactive and proactive aggression, callous-unemotional traits, default mode network, amygdala, functional connectivity

[A bioRxiv preprint version of this manuscript has been updated on April 6th 2019]

3 Study B

DIFFERENT WHOLE-BRAIN FUNCTIONAL CONNECTIVITY CORRELATES OF REACTIVE-PROACTIVE AGGRESSION AND CALLOUS-UNEMOTIONAL TRAITS IN DISRUPTIVE CHILDREN AND ADOLESCENTS

Julia E Werhahn^{1,2}, Susanna Mohl¹, David Willinger^{1,2}, Lukasz Smigielski^{1,2}, Alexander Roth¹, Jilly Naaijen^{3,4}, Leandra M Mulder^{3,4}, Jeffrey C Glennon³, Pieter J Hoekstra⁵, Andrea Dietrich⁵, Renee Kleine Deters⁵, Pascal M Aggensteiner⁶, Nathalie E Holz⁶, Sarah Baumeister⁶, Tobias Banaschewski⁶, Melanie C Saam⁷, Ulrike M E Schulze⁷, David J Lythgoe⁸, Arjun Sethi⁸, Michael Craig⁸, Mathilde Mastroianni⁹, Ilyas Sagar-Ouriaghli⁹, Paramala J Santosh⁹, Mireia Rosa¹⁰, Nuria Bargallo¹¹, Josefina Castro-Fornieles¹², Celso Arango¹³, Maria J Penzol¹³, Marcel P Zwiers⁴, Barbara Franke^{14,15}, Jan K Buitelaar^{3,16}, Susanne Walitza^{1,2, 17}, Daniel Brandeis^{1,2,6, 17}

¹Department of Child and Adolescent Psychiatry and Psychotherapy, University Hospital of Psychiatry Zurich, University of Zurich, Zurich, Switzerland

²Neuroscience Center Zurich, University and ETH Zurich, Zurich, Switzerland

³Radboud University Medical Center, Donders Institute for Brain, Cognition and Behavior, Department of Cognitive Neuroscience, Nijmegen, The Netherlands

⁴Radboud University, Donders Institute for Brain, Cognition and Behavior, Centre for Cognitive Neuroimaging, Nijmegen, The Netherlands

⁵University of Groningen, University Medical Center Groningen, Department of Child and Adolescent Psychiatry, Groningen, The Netherlands

⁶Department of Child and Adolescent Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim/Heidelberg University, Mannheim, Germany

⁷Department of Child and Adolescent Psychiatry/Psychotherapy, University Hospital, University of Ulm, Germany

⁸Department of Neuroimaging, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

⁹Department of Child Psychiatry, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

¹⁰Child and Adolescent Psychiatry Department, Hospital Clinic of Barcelona, IDIBAPS, Barcelona, Spain

¹¹Clinic Image Diagnostic Center (CDIC), Hospital Clinic of Barcelona; Magnetic Resonance Image Core Facility, IDIBAPS, Barcelona, Spain

¹²Child and Adolescent Psychiatry and Psychology Department, 2017SGR881, Institute Clinic of Neurosciences, Hospital Clinic of Barcelona, CIBERSAM, IDIBAPS, Department of Medicine, University of Barcelona, Barcelona, Spain

¹³Child and Adolescent Psychiatry Department, Hospital General Universitario Gregorio Marañón School of Medicine, Universidad Complutense, IiSGM, CIBERSAM, Madrid, Spain

¹⁴Radboud University Medical Center, Donders Institute for Brain, Cognition and Behavior, Department of Human Genetics, Nijmegen, The Netherlands

¹⁵Radboud University Medical Center, Donders Institute for Brain, Cognition and Behavior, Department of Psychiatry, Nijmegen, The Netherlands

¹⁶Karakter Child and Adolescent Psychiatry University Center, Nijmegen, The Netherlands

¹⁷Zurich Center for Integrative Human Physiology, University of Zurich, Zurich, Switzerland

Short title: Whole-brain aggression-related connectivity

Keywords: reactive aggression; proactive aggression; callous-unemotional traits, fMRI; resting state; functional connectivity; anxiety; ADHD; children; adolescents

4 General Discussion

In this dissertation project, we investigated connectivity alterations during rs-fMRI in children and adolescents with clinical levels of aggression, and mostly with diagnoses of CD and/or ODD, compared to typically developing healthy controls. Given the limited knowledge of neural correlates underlying partly distinct manifestations of aggressive behavior at rest, particularly RA and PA symptomatology, we extended previous studies by elucidating rsFC related to these aggression subtypes along with CU traits. Both seed-based analysis (study A) and voxel-to-voxel approaches (study B) revealed rsFC alterations in cases compared to healthy controls and, most importantly, predominantly distinct aggression subtype-specific patterns. Furthermore, these results emphasized the importance of considering also clinical symptoms beyond aggression, like comorbid ADHD and anxiety symptoms. After the discussion of our findings including potential directions for future studies, implications for the clinical practice regarding the eligibility of rsFC patterns as a biomarker for aggression-related behaviors are outlined.

4.1 Resting state fMRI in aggressive children and adolescents – main findings and future directions

Investigations of rs-fMRI, which detect correlated low-frequency time courses of brain activity during periods of rest, have the advantages of being non-invasive, providing a good spatial resolution (Huettel et al., 2009), and being independent of higher levels of cooperation (Smitha et al., 2017) which facilitates rs-fMRI data collection in our population of aggressive children and adolescents. Yet, as our cases with externalizing disorders including comorbid ADHD symptoms showed higher levels of motion compared to the healthy controls, we used a threshold for excessive motion

recently applied (Pruim, Mennes, Buitelaar, & Beckmann, 2015) also in rs-fMRI analyses in adolescents with ADHD (von Rhein et al., 2016). Above that, we conducted sensitivity analyses controlling for effects of this motion parameter, which yielded comparable statistics in our case-control differences in rsFC. Despite growing research on behavioral and psychophysiological levels, neural studies of rs-fMRI in aggressive children and adolescents have widely neglected the differentiation between distinct manifestations of aggressive behavior, particularly RA and PA symptomatology along with comorbid ADHD and anxiety symptoms. Besides addressing this gap in literature, the present PhD project further extended previous research by not exclusively including male adolescents with CD but also females, children, and individuals with ODD and/or aggressive behavior in the clinical range, and providing a considerably larger sample size ($N = 207$) including 118 cases.

Both our seed and voxel-based rsFC findings showed that RA and PA symptoms related to predominantly distinct connectivity patterns. However, we found a converging left amygdala seed-based pattern with precuneal areas and similar voxel-wise rsFC changes including superior parietal regions related to both RA and PA symptoms. Moreover, RA and PA scores were highly correlated in our sample in line with the literature (Cima et al., 2013). These neural and behavioral similarities involving the amygdala as core region implicated in aggression might challenge the proposed distinction between RA and PA aggression subtypes. However, this amygdala seed-based pattern extended to different cingulate and occipital areas for PA and RA scores, respectively, RA scores only related to further seed-based patterns, and both forms of aggression correlated with distinct voxel-wise connectivity alterations. These RA and PA-specific results thus corroborate behavioral studies reporting differential behavioral correlates for RA and PA symptomatology (Fite et al., 2008, 2010; Marsee et al., 2011; Marsee & Frick, 2007).

Additionally, both studies revealed that the left anterior insula is implicated in CU traits-related rsFC alterations. Adults with psychopathic traits exhibited rsFC alterations including the anterior insula (Philippi et al., 2015), which might point to developmental aspects for this brain area given

that CU traits in childhood and psychopathic traits in adulthood correlate on a behavioral level (Frick et al., 2013). Moreover, some of the brain regions involved in the CU traits-related connectivity patterns reported in our studies have been linked to rsFC alterations in adults with psychopathic traits, such as the precuneus (Motzkin et al., 2011; Philippi et al., 2015) and central and angular gyrus (Espinoza et al., 2018). Furthermore, our results showed CU trait-specific rsFC alterations beyond (para-) limbic areas in frontal, parietal, cingulate, and cerebellar regions, which points in the direction of a recent review that reported broader brain areas related to psychopathy in adulthood (Griffiths & Jalava, 2017). Additionally, in both aggressive children and adolescents of our sample and in adult violent offenders of a recent study (Siep et al., 2018) RA and PA scores related to left amygdala seed-based rsFC. Thus, longitudinal studies could identify developmental trajectories by observing rsFC in these specific brain regions over time. These investigations might clarify whether these deviations reflect early onset aggressive symptoms that persist into adulthood or whether they predict future aggression-related outcomes. In both cases, designing early neurobiological interventions specifically targeted to alter connectivity patterns including these areas might be promising, as they could eventually improve behavioral outcomes. For instance, the anterior insula has been shown to be implicated during self-regulation in real-time fMRI neurofeedback (Emmert et al., 2016) and has already been targeted in adult psychopaths (Sitaram et al., 2014). Moreover, as changes in electrodermal activity were shown to involve activity in brain regions including the insular cortices (Nagai, Critchley, Featherstone, Trimble, & Dolan, 2004), arousal-biofeedback training (Schoenberg & David, 2014) may be promising and is thus part of our Aggrosotype project.

Interestingly, we found some regions implicated in both RA/PA symptoms and CU traits-related rsFC. For instance, seed-based analyses showed that clusters including precentral gyrus and precuneus were implicated in both RA/PA symptoms-related patterns and those associated with CU subdimensions. Additionally, voxel-wise patterns related to RA and PA symptoms included parietal regions such as the angular gyrus, previously linked to CU traits-related activity alterations (Anderson et al., 2017). Yet,

voxel-wise and seed-based RA and PA symptoms-specific rsFC patterns in the present dissertation differed from those associated with CU traits. This contradicts previous behavioral reports of a correlation between RA and PA symptoms with CU traits (Urban et al., 2018) and the correlation of PA symptoms and CU traits in our sample according to behavioral measures. The RA and PA symptoms-specific rsFC results differing from the CU traits-related findings thus add further evidence to the knowledge of distinct underlying neural correlates for these manifestations of aggression.

Additionally, our findings strongly suggest that research (and clinical practice) should consider comorbid symptoms in disruptive children and adolescents given that these comorbidities exert suppressor effects on numerous rsFC patterns. Thereby, the reduced seed-based limbic-frontal rsFC in cases compared to controls depended on the additional control for ADHD symptoms, and the voxel-based connectivity alterations including the medial frontal cortex vanished without the additional control for anxiety problems. Above that, most voxel-based rsFC subtype-specific alterations within cases could only be identified when both ADHD and anxiety symptoms were considered. This provides support for previous neural studies regarding ADHD symptoms in rsFC within the DMN in male adolescents with CD (Broulidakis et al., 2016), and regarding anxiety levels in rsFC (Motzkin et al., 2011) and task-based fMRI (Sethi, Sarkar, et al., 2018) in psychopathic adults. In our studies, we extended these findings to brain regions like the SN and to further aggression subtypes including RA and PA symptomatology. The behavioral correlations, particularly within cases, provide further support for correlations between anxiety and both RA and PA symptoms (Fite et al., 2014; Fung et al., 2015) or with RA symptoms particularly (Marsee et al., 2008; Vitaro et al., 2002) in previous behavioral studies. Our rsFC results also corroborate behavioral research emphasizing the crucial role of both ADHD (Fonagy & Luyten, 2017) and anxiety symptoms (Frick, 2012; Frick et al., 2013) in disruptive children and adolescents. A recent longitudinal study documented a negative association between anxiety and CU-related traits at age six and, beyond that, anxiety levels predicting CU-related traits six years later (Bamvita et al., 2017). Another dimensional severity-based approach found that youths with both

high CU traits and high compared to low levels of anxiety differ in factors, such as a more severe trauma history in childhood and ADHD symptoms (Cecil, McCrory, Barker, Guiney, & Viding, 2018). CU traits combined with high levels of anxiety further related to distinct characteristics like increased aggression and conduct problems (Fanti et al., 2013) as in our sample, along with reduced self-reported cognitive empathy (Kahn, Frick, Golmaryami, & Marsee, 2017), increased antisocial behavior and peer problems (Humayun, Kahn, Frick, & Viding, 2014). On the contrary, increased CU traits in youths with low anxiety levels related to reduced emotion recognition skills (Dadds et al., 2017), emotional intelligence (Kahn, Ermer, Salovey, & Kiehl, 2016), perceived positive outcomes of antisocial behavior and higher participation in these behavior (Guelker et al., 2014), and emotion-related deficits (Kimonis et al., 2012). Given this growing body of literature on a behavioral level, it is somehow surprising that rsFC analysis to date neglected the role of comorbid anxiety symptoms and only few considered co-occurring ADHD symptoms.

Furthermore, both analysis approaches showed that cases and controls differed only in few rsFC patterns leaving limited room for interpretation, as the differences only survived the correction for site but not for further covariates including sex, IQ, handedness, and medication. In contrast, we observed several dimensional aggression subtype-specific results within cases derived from both seed and voxel-based analyses, which broadly survived additional corrections. This point to the heterogeneity within aggression-related disorders and, above that, support rather a dimensional than categorical approach as increasingly promoted in recent years (Blair et al., 2016; Wesselhoeft et al., 2018).

In contrast to the majority of previous rsFC-studies, our projects included not only male but also female participants. The distribution of sex in our sample reflected prevalence estimates, for instance regarding CD (Loeber, Burke, & Pardini, 2009). Yet, from a statistical point, the smaller number of female patients with aggressive behavior in the clinical range ($N = 19$) compared to male cases ($N = 99$) and, above that, recruited from different sites, makes it statistically difficult to draw reliable conclusions on sex-related effects. As recently documented, adolescents with CD showed sex-

related differences in spontaneous brain activity during rest including left ITG/MTG (Cao et al., 2018). In our sample of aggressive boys and girls, comparable temporal regions exhibited CU traits-related changes in voxel-wise patterns, which survived the post hoc control for additional covariates including sex. Nevertheless, we included this variable as a covariate and conducted corresponding sensitivity analyses. However, future studies could elucidate sex-related differences in aggression subtype-specific rsFC.

Regarding the applied statistical approaches, we decided to use a seed-based approach with whole-brain voxel clusters on the one hand and voxel-to-voxel graph-theory-based unrestricted whole-brain analyses on the other hand. The common seed-based approach (Aghajani et al., 2016, 2017; Pujol et al., 2012; Uytun et al., 2016) provides us with information of brain areas functionally connected with a seed and a direct interpretation of the results (Mak et al., 2017). While ICA enables to identify spatially distinct RSNs (Beckmann, DeLuca, Devlin, & Smith, 2005), seed-based analysis do not require to manually select important components and to distinguish between noise and physiological signal (Lee et al., 2013). Moreover, results of ICA and seed-based analysis seem comparable (Rosazza, Minati, Ghielmetti, Mandelli, & Bruzzone, 2012). Our second study included the calculation of both global (ICC) and local (ILC) voxel-wise connectivity changes. ICC has been frequently applied in recent rsFC studies (Browndyke et al., 2018; Cassady et al., 2018; Van Ombergen et al., 2017; Vatansever et al., 2017; Walpola et al., 2017) and is used to calculate the global connectivity strength and determine degree or network centrality (Martuzzi et al., 2011). Among the advantages of ICC is the circumvention of the need for a priori definitions of ROIs, as required for seed-based analysis. ILC integrates spatial correlations to calculate average local connectivity patterns and provides us with an independence of image resolution and no need for predefining the neighborhood, as necessary for other local coherence approaches such as regional homogeneity (Deshpande et al., 2009). Given our considerable larger data set, it would be of great interest to determine rsFC patterns by using classification methods (Abós et al., 2017), as recently applied to identify levels of psychopathology in children and adolescents (Sato et al., 2018) and on a structural level to distinguish

male adolescents with conduct disorder from healthy controls (Zhang et al., 2018). This approach could enable to elucidate connectivity patterns related to aggression subtype-specific behavioral measures, which might provide further support for an aggression subtype-specific approach.

Furthermore, one has to keep in mind that MRI-derived data provides a good spatial resolution, but data collected with other imaging methods like the electroencephalogram (EEG) provides a higher temporal resolution (Huettel et al., 2009). Thus, in order to widen the knowledge of aggression-related neural correlates with both a good location of the signal and more precise information regarding oscillations of brain activity (Huettel et al., 2009; Meyer-Lindenberg, 2010), future studies could investigate connectivity alterations during task-based fMRI paradigms combined with EEG measures.

For this dissertation, data collection took place at nine sites across Europe using differing scanners with partly diverging scan parameters. Implementing a multi-echo resting state sequence at all participating sites would have further improved the differentiation between BOLD and artifact signal (Kundu et al., 2017). In order to control for site effects, we added a dummy-coded site covariate in all rsFC analyses, as sensitivity analyses revealed a significant effect of site on some rsFC patterns. However, we collected data from travelling heads at each site and defined consistent inclusion and exclusion criteria, which included standard clinical measures. Above that, activation variability due to site specific variations has been shown to be small in other studies (Yendiki et al., 2010), and recent multi-centric rs-fMRI-studies successfully analyzed data from multiple sites (Biswal et al., 2010; Chen et al., 2016; Jovicich et al., 2016; Tomasi & Volkow, 2012). Moreover, the larger sample size and multi-center design might improve reliability and generalizability of our results. Despite these advantages, we did not explicitly account for potential effects based on cultural differences. Yet, there is evidence for systematic mechanisms to affect the assessment of child problem behavior vary across cultural minorities, as members of some cultures might tend to report aggressive behavior to a lesser extend (Eisner & Ribeaud, 2007). Furthermore, a recent paper points to the importance of considering the role of social norms

regarding neurobiological differences of aggression (Porter, 2018). Consequently, future studies could address cultural differences by systematically document cultural origins of the participants in order to control potential culture-related differences and increase generalizability of aggression-related correlates.

With respect to interpretation of the observed rsFC results, there are some limitations worth mentioning. The number of rs-fMRI studies in aggressive children and adolescents is limited, mostly restricted to male adolescents with CD exclusively, characterized by differing methodological approaches, and with respect to RA and PA subtypes and comorbid anxiety symptoms non-existent. This made it considerably difficult to embed and interpret the observed rsFC patterns in the present PhD project. Noteworthy, some of these previous rs-fMRI studies used the same sample of male adolescents with CD and healthy controls five times (Lu, Zhou, Wang, et al., 2017; Lu, Zhou, Zhang, et al., 2017; Lu et al., 2015; Zhou et al., 2016, 2015). This might elucidate the knowledge of implicated brain areas revealed by differing methodological approaches though hardly enables transferring these results to a broader population of patients with aggression-related disorders. Therefore, we tentatively linked altered functioning during rest observed in our sample of aggressive children and adolescents to results reported in similar brain regions in task-based functional and structural MRI studies.

Moreover, it is important to note that rs-fMRI data in general does not allow for conclusions on brain activity or connectivity in response to stimuli or during a specific task to be drawn. Compared to task-based fMRI, though, rs-fMRI has many advantages. For instance, it provides us with an improved signal-to-noise-ratio, enables investigating a considerable number of brain functions by applying different analyses approaches while using the same data set, and is independent of higher levels of cooperation as needed for task-based designs (Smitha et al., 2017). Moreover, RSNs like the DMN (Andrews-Hanna et al., 2010; Fox et al., 2015; Raichle et al., 2001) can be identified during rs-fMRI across all age groups (Mak et al., 2017) including children (Thomason et al., 2011), and rsFC patterns were shown to exhibit

moderate to high test-retest reliability across subjects and sessions (van Dijk et al., 2010).

Yet, evidence from aggression subtype-specific structural and task-based functional MRI is inevitable to draw further conclusions that might eventually effect diagnostic and treatment approaches. To date, these interventions designed to treat aggression-related disorders in children and adolescents show small effect sizes (Bakker et al., 2017) and seem to be less effective in the presence of CU traits (Wilkinson et al., 2016). There are suggestions that children and adolescents with high levels of CU traits might profit from interventions that consider environmental factors such as parenting practices (Fonagy & Luyten, 2017) and, broadly speaking, from an individualized treatment according to the distinct manifestation of aggression (Frick, 2016). For instance, there is evidence for a better response in children with conduct problems and additional CU traits to emotion-recognition-training (Dadds, Cauchi, Wimalaweera, Hawes, & Brennan, 2012). The results documented in the present PhD project encourage transferring an aggression subtype-specific approach to the clinical practice. Thus, diagnostic and treatment approaches might not only profit from considering CU-traits (Umbach et al., 2015; Wilkinson et al., 2016) but also to RA and PA along with comorbid ADHD and anxiety symptoms.

Taken together, our results shed light on aberrant rsFC patterns in aggressive children and youths compared to healthy controls and on the essential role of distinct manifestations of aggressive behaviors including comorbid ADHD and anxiety symptoms within cases. Besides several limitations, such as possibly lowered study power due to partly deviating scan parameters, the reasonable larger sample size and multi-center design are among the strengths of our studies, as they may increase reliability and generalizability of the reported results. The rsFC analyses conducted in the present PhD project and the resulting aggression-related connectivity patterns add to further characterizing and disentangling task-independent neural correlates of different aggressive behaviors in children and adolescents. Future studies could reveal aggression-related connectivity patterns at a higher temporal resolution by applying EEG measurements

during task-based designs. Longitudinal designs might clarify the role of aggression subtype-specific neural correlates such as amygdala and insula over time, with promising implications for early interventions.

4.2 Clinical implications – functional connectivity alterations during rest as a biomarker in disruptive children and adolescents?

The present PhD project investigated aggression-related connectivity aberrations during rs-fMRI in children and adolescents. Our main aim was to understand neural alterations related to distinct forms of aggression, particularly RA and PA symptomatology along with CU traits. Both studies revealed aggression subtype-specific rsFC patterns, thereby extended the knowledge of aggression-related neural networks to distinct aggression subtypes. Beside these new insights, our results further underlined the crucial role of considering ADHD and anxiety symptoms, frequently comorbid in aggression-related disorders though widely disregarded to date.

Neuroimaging studies could extend the knowledge of underlying mechanisms, design tailored treatment, and investigate treatment effects (Umbach et al., 2015). Hence, clinicians would be able to not only tailor treatment accordingly but also monitor progress and evaluate the effects of interventions (Puzzo et al., 2016). Using rs-fMRI in particular, patients with psychiatric diagnoses, such as major depressive disorder, autism, or ADHD, have been distinguished from controls (Lee et al., 2013). Given advantages like the relatively ease of acquiring rs-fMRI data, rs-fMRI-derived patterns might be a promising future biomarker and additional diagnostic tool for psychiatric disorders (Woodward & Cascio, 2015), also in children (Thomason et al., 2011). Using research of the Human Connectome Project and further large databases could facilitate linking behavioral characteristics to rsFC patterns (Woodward & Cascio, 2015). The potential use of rsFC alterations as a biomarker in children and adolescents with

aggressive behavior in the clinical range, accompanied by several challenges and limitations, will be outlined in the following.

Amid the heterogeneity of aggression-related disorders with diverging implications for treatment (Byrd et al., 2014; Umbach et al., 2015; Wilkinson et al., 2016), the need for investigating aggression subtypes is evident. To address the variety in aggression-related disorders, the Research Domain Criteria (RDoC) approach is of particular interest, as it integrates different domains (Fonagy & Luyten, 2017) and considers mental disorders in the light of dysfunctions in neural circuits (Insel et al., 2010). Moreover, given the growing body of evidence of dysfunctional neural systems in aggression-related disorders including limbic, paralimbic, and frontal areas associated with cognitive impairments (Blair et al., 2016), the RDoC approach might be useful. Thus, syndromes could be characterized based on the identification of underlying trajectories on different levels including neuroimaging, which would optimize diagnostic and treatment as defined by precision psychiatry (Silbersweig & Loscalzo, 2017). A recent paper integrated theoretical approaches and empirical findings regarding aggression and conduct problems from the RDoC perspective, and underlines the importance of considering biological, psychological, and environmental factors (Fonagy & Luyten, 2017). For instance, the authors identified impaired empathic responses and emotional perspective taking as relevant etiological factors, and two main clusters of children and adolescents with high anxious, reactive aggressive behaviors on the one side and impaired affective mentalizing, instrumental form of aggression on the other side. These RDoC-derived findings corroborate our observed distinct RA and PA symptoms-related rsFC patterns along with the crucial role of considering co-occurring anxiety symptoms.

The RDoC approach, though, is accompanied by several challenges. For instance, one important challenge is to consider the differing outcomes related to the same diagnosis, as reflected by the heterogeneity within conduct disorder. Therefore, distinct factors can lead to similar rsFC profiles (Lilienfeld, 2014). In CD, there is evidence for genetic and environmental factors and gene-environment interplays (Salvatore & Dick, 2016). For instance, trauma history has been shown to play an important role in

differentiating CU youths with high and low anxiety levels (Cecil et al., 2018), and childhood family adversity including poor parental monitoring seems to predict structural and functional alterations in young adults in brain areas like the amygdala (Holz et al., 2016). Regarding rs-fMRI, early-life stress exposure in young children seems related to higher regional homogeneity during rs-fMRI in the left lateral frontal cortex (Demir-Lira et al., 2016). Therefore, a holistic and comprehensive understanding of rsFC patterns as possible biomarker for aggression-related disorders seems inevitable. Thus, we need to consider not only biological predispositions but also whether these predispositions effectively resulted in disruptive behavior (Lilienfeld, 2014). This might be of high relevance to inform subsequent targeted therapeutic interventions accordingly. In sum, the RDoC approach could enable to capture and determine the heterogeneity of psychiatric disorders including clinical manifestations of aggression-related disorders (Fonagy & Luyten, 2017). Given its limitations, particularly regarding the equifinality of differing developmental pathways with diverging implications for treatment, this framework might be of rather supplemental utility (Lilienfeld, 2014).

Furthermore, information about developmental trajectories derived from longitudinal studies might contribute to determining specific rsFC patterns as biomarker for aggression-related disorders and their distinct subtypes and manifestations in childhood and adolescence. At the same time, possible maturation-related changes in rsFC (Thomason et al., 2011; van Dijk et al., 2010) would have been taken into account, for instance in prefrontal and precuneal areas (Chen et al., 2015).

Moreover, the differing analysis methods in the present and in previous studies need evaluation regarding their efficacy in identifying psychiatric symptomatologies (Lee et al., 2013). It is noteworthy though that the different methodological approaches applied in our studies revealed partly overlapping brain regions implicated in rsFC alterations. For instance, both analysis methods showed PA-related connectivity alterations in precuneal areas. The precuneus has been associated with cool executive functioning, which describes behaviors such as self-regulation (Diamond, 2013). Interestingly, a recent narrative review reported reduced precuneal activity

linked to ODD/CD (Noordermeer et al., 2016). Moreover, a recent meta-analytic study related trait aggression and executive functioning to aberrant brain activity including the precuneus (Wong et al., 2018). Beside the PA-related precuneal patterns, case-control differences including the frontal pole and CU traits-related rsFC alterations including the left insula within cases resulted from both seed and voxel-based analysis. Therefore, these areas might be of high relevance for subtyping aggressive behavior on a neural level.

Finally, in order to use rsFC patterns as a predictive biomarker in aggression-related disorders and their distinct manifestations, future research is needed to replicate our findings particularly regarding RA and PA aggression subtypes and comorbid anxiety symptoms, as they have been investigated in this PhD project for the first time in the field of rs-fMRI. Taken together, the aggression subtype-specific rsFC patterns documented here showed that differing analysis approaches – revealing similar subtype-specific areas implicated like the left anterior insula and the precuneus – can enable to disentangle distinct manifestations of aggressive behaviors. Thus, these patterns might be promising neuroimaging biomarkers in children and adolescents with disruptive behavior.

4.3 Conclusions

In this PhD project, we applied two analysis approaches to investigate aggression-related aberrations of brain function at rest in a rs-fMRI sequence acquired from a relatively larger sample ($N > 200$) of children and adolescents recruited in the framework of the European multi-center projects Aggresstotype/MATRICS. First, cases showed both seed and voxel-based connectivity alterations in brain areas including the frontal pole compared to healthy controls. These group differences partly depended on the additional control of co-occurring ADHD or anxiety symptoms respectively. Second, both studies revealed predominantly distinct RA and PA symptoms-related rsFC alterations in brain areas including the precuneus along with CU traits-related patterns involving the left anterior insula within cases. Third, particularly the observed aggression subtype-

specific voxel-wise results emphasized the importance of additional considering both comorbid ADHD symptoms and anxiety levels.

This dissertation points to the importance of differentiating between RA and PA symptoms and CU traits in children and adolescents with aggressive behavior in the clinical range on a neural level. Moreover, considering comorbid levels of ADHD and anxiety seems inevitable. Our results add to the knowledge of neural characteristics implicated in different aggression-related behaviors and their specific clinical manifestations. The present PhD project further encourages a careful assessment of distinct manifestations of aggressive behaviors and comorbid symptoms following clinical standards and guidelines as well as using additional research instruments. Our results might be promising predictive biomarkers for specific forms of aggression in children and adolescents. Above that, they may eventually inform improved diagnostic and treatment of aggression-related disorders thus tailored accordingly, as already implemented in the individualized arousal-biofeedback training in our Aggessotype project.

Abbreviations

ADHD	attention deficit/hyperactivity disorder
amPFC	anterior medial prefrontal cortex
BOLD	blood-oxygen-level dependent
CD	conduct disorder
CU	callous-unemotional
DMN	default mode network
DSM-5	Diagnostic and Statistical Manual of Mental Disorders 5
FDR	false discovery rate (correction)
fMRI	functional magnetic resonance imaging
FWE	family-wise error (correction)
GLM	generalized linear model
ICC	intrinsic connectivity contrast
ILC	integrated local correlation
MNI	Montreal Neurological Institute
MRI	magnetic resonance imaging
ODD	oppositional defiant disorder
OFC	orbitofrontal cortex
PA	proactive aggression
PCC	posterior cingulate cortex
RA	reactive aggression
ROI	region of interest
RDoC	research domain criteria
rsFC	resting state functional connectivity
rs-fMRI	resting state functional magnetic resonance imaging
RSN	resting state network

SN salience network

References

- Abós, A., Baggio, H. C., Segura, B., García-Díaz, A. I., Compta, Y., Martí, M. J., ... Junqué, C. (2017). Discriminating cognitive status in Parkinson's disease through functional connectomics and machine learning. *Scientific Reports*, 7(November 2016), 1–13.
- Achenbach, T. M. (1991). *Integrative guide for the 1991 CBCL/4-18, YSR, and TRF profiles*. Burlington, VT: University of Vermont, Department of Psychiatry.
- Aghajani, M., Colins, O. F., Klapwijk, E. T., Veer, I. M., Andershed, H., Popma, A., ... Vermeiren, R. R. J. M. (2016). Dissociable relations between amygdala subregional networks and psychopathy trait dimensions in conduct-disordered juvenile offenders. *Human Brain Mapping*, 37(11), 4017–4033.
- Aghajani, M., Klapwijk, E. T., van der Wee, N. J., Veer, I. M., Rombouts, S. A. R. B., Boon, A. E., ... Colins, O. F. (2017). Disorganized Amygdala Networks in Conduct-Disordered Juvenile Offenders With Callous-Unemotional Traits. *Biological Psychiatry*, 82(4), 283–293.
- Alegria, A. A., Radua, J., & Rubia, K. (2016). Meta-analysis of fmri studies of disruptive behavior disorders. *American Journal of Psychiatry*, 173(11), 1119–1130.
- Allen, E. A., Erhardt, E. B., Damaraju, E., Gruner, W., Segall, J. M., Silva, R. F., ... Calhoun, V. D. (2011). A baseline for the multivariate comparison of resting-state networks. *Frontiers in Systems Neuroscience*, 5(February), 2.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders: DSM-5*. Washington, DC: American Psychiatric Association.
- Anderson, N. E., Steele, V. R., Maurer, J. M., Rao, V., Koenigs, M. R., Decety, J., ... Kiehl, K. A. (2017). Differentiating emotional processing and attention in psychopathy with functional neuroimaging. *Cognitive, Affective and Behavioral Neuroscience*, 17(3), 491–515.
- Andersson, J. L. R., Hutton, C., Ashburner, J., Turner, R., & Friston, K. (2001).

- Modeling Geometric Deformations in EPI Time Series. *NeuroImage*, 13(5), 903–919.
- Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R., & Buckner, R. L. (2010). Functional-Anatomic Fractionation of the Brain's Default Network. *Neuron*, 65(4), 550–562.
- Andrews-Hanna, J. R., Smallwood, J., & Spreng, R. N. (2014). The default network and self-generated thought: Component processes, dynamic control, and clinical relevance. *Annals of the New York Academy of Sciences*, 1316(1), 29–52.
- Bakker, M. J., Greven, C. U., Buitelaar, J. K., & Glennon, J. C. (2017). Practitioner Review: Psychological treatments for children and adolescents with conduct disorder problems - a systematic review and meta-analysis. *Journal of Child Psychology and Psychiatry*, 1, 4–18.
- Bamvita, J. M., Larm, P., Checknita, D., Vitaro, F., Tremblay, R. E., Côté, G., & Hodgins, S. (2017). Childhood predictors of adult psychopathy scores among males followed from age 6 to 33. *Journal of Criminal Justice*, 53(September), 55–65.
- Barch, D. M. (2017). Resting-state functional connectivity in the human connectome project: Current status and relevance to understanding psychopathology. *Harvard Review of Psychiatry*, 25(5), 209–217.
- Beckmann, C. F., DeLuca, M., Devlin, J. T., & Smith, S. M. (2005). Investigations into resting-state connectivity using independent component analysis. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1457), 1001–1013.
- Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, 37(1), 90–101.
- Biswal, B. B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S. M., ... Milham, M. P. (2010). Toward discovery science of human brain function. *Proceedings of the National Academy of Sciences*, 107(10), 4734–4739.
- Biswal, B., Yetkin, F., Haughton, V., & Hyde, J. (1995). Functional connectivity in the motor cortex of resting human brain using. *Magn Reson Med*, 34(9), 537–541.
- Blair, R. J. R. (2001). Neurocognitive models of aggression, the antisocial personality disorders, and psychopathy. *Journal of Neurology, Neurosurgery & Psychiatry*, 71(6), 727–731.
- Blair, R. J. R. (2004). The roles of orbital frontal cortex in the modulation of antisocial behavior. *Brain and Cognition*, 55(1), 198–208.

- Blair, R. J. R. (2010). Psychopathy, frustration, and reactive aggression: the role of ventromedial prefrontal cortex. *British Journal of Psychology (London, England : 1953)*, 101(Pt 3), 383–399.
- Blair, R. J. R. (2016). The neurobiology of impulsive aggression. *Psychiatric Clinics of North America*, 26(1), 4–9.
- Blair, R. J. R. (2017). Emotion-based learning systems and the development of morality. *Cognition*, 167, 38–45.
- Blair, R. J. R., Leibenluft, E., & Pine, D. S. (2014). Conduct disorder and callous-unemotional traits in youth. *The New England Journal of Medicine*, 371(23), 2207–2216.
- Blair, R. J. R., Veroude, K., & Buitelaar, J. K. (2016). Neuro-cognitive system dysfunction and symptom sets: A review of fMRI studies in youth with conduct problems. *Neuroscience and Biobehavioral Reviews*.
- Blais, J., Solodukhin, E., & Forth, A. E. (2014). A Meta-Analysis Exploring the Relationship Between Psychopathy and Instrumental Versus Reactive Violence. *Criminal Justice and Behavior*, 41(7), 797–821.
- Boccia, M., Dacquino, C., Piccardi, L., Cordellieri, P., Guariglia, C., Ferlazzo, F., ... Giannini, A. M. (2017). Neural foundation of human moral reasoning: an ALE meta-analysis about the role of personal perspective. *Brain Imaging and Behavior*, 11(1), 278–292.
- Bordin, I. A., Rocha, M. M., Paula, C. S., Teixeira, M. C. T. V., Achenbach, T. M., Rescorla, L. A., & Silveiras, E. F. M. (2013). Child Behavior Checklist (CBCL), Youth Self-Report (YSR) and Teacher's Report Form (TRF): an overview of the development of the original and Brazilian versions. *Cadernos de Saúde Pública*, 29(1), 13–28.
- Broulidakis, M. J., Fairchild, G., Sully, K., Blumensath, T., Darekar, A., & Sonuga-Barke, E. J. S. (2016). Reduced Default Mode Connectivity in Adolescents With Conduct Disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*.
- Brovelli, A., Nazarian, B., Meunier, M., & Boussaoud, D. (2011). Differential roles of caudate nucleus and putamen during instrumental learning. *NeuroImage*, 57(4), 1580–1590.
- Browndyke, J. N., Berger, M., Smith, P. J., Harshbarger, T. B., Monge, Z. A., Panchal, V., ... Mathew, J. P. (2018). Task-related changes in degree centrality and local coherence of the posterior cingulate cortex after major cardiac surgery in older adults. *Human Brain Mapping*, 39(2), 985–1003.
- Buss, D. M., & Shackelford, T. K. (1997). Human aggression in evolutionary

- psychological perspective. *Clinical Psychology Review*, 17(6), 605–619.
- Byrd, A. L., Loeber, R., & Pardini, D. A. (2014). Antisocial Behavior, Psychopathic Features and Abnormalities in Reward and Punishment Processing in Youth. *Clinical Child and Family Psychology Review*, 17(2), 125–156.
- Bzdok, D., Schilbach, L., Vogeley, K., Schneider, K., Laird, A. R., Langner, R., & Eickhoff, S. B. (2012). Parsing the neural correlates of moral cognition: ALE meta-analysis on morality, theory of mind, and empathy. *Brain Structure and Function*, 217(4), 783–796.
- Calhoun, V. D., Wager, T. D., Krishnan, A., Rosch, K. S., Seymour, K. E., Nebel, M. B., ... Kiehl, K. (2017). The impact of T1 versus EPI spatial normalization templates for fMRI data analyses. *Human Brain Mapping*, 38(11), 5331–5342.
- Cao, W., Sun, X., Dong, D., Yao, S., & Huang, B. (2018). Sex differences in spontaneous brain activity in adolescents with conduct disorder. *Frontiers in Psychology*, 9(AUG), 1–9.
- Card, N. A., & Little, T. D. (2006). Proactive and reactive aggression in childhood and adolescence: A meta-analysis of differential relations with psychosocial adjustment. *International Journal of Behavioral Development*, 30(5), 466–480.
- Cassady, K., Ruitenberg, M., Koppelmans, V., Reuter-Lorenz, P., De Dios, Y., Gadd, N., ... Seidler, R. (2018). Neural predictors of sensorimotor adaptation rate and savings. *Human Brain Mapping*, 39(4), 1516–1531.
- Cavanna, A. E. (2007). The precuneus and consciousness. *CNS Spectrums*, 12(7), 545–552.
- Cecil, C. A. M., McCrory, E. J., Barker, E. D., Guiney, J., & Viding, E. (2018). Characterising youth with callous-unemotional traits and concurrent anxiety: evidence for a high-risk clinical group. *European Child and Adolescent Psychiatry*, 27(7), 885–898.
- Chai, X. J., Nieto-Castanon, A., Öngür, D., & Whitfield-Gabrieli, S. (2012). Anticorrelations in resting state networks without global signal regression. *Neuroimage*, 59(3), 1190–1199.
- Chen, C., Zhou, J., Liu, C., Witt, K., Zhang, Y., Jing, B., ... Li, L. (2015). Regional Homogeneity of Resting-State Brain Abnormalities in Violent Juvenile Offenders: A Biomarker of Brain Immaturity? *Journal of Neuropsychiatry and Clinical Neurosciences*, 27(1), 27–32.
- Chen, H., Duan, X., Liu, F., Lu, F., Ma, X., Zhang, Y., ... Chen, H. (2016).

- Multivariate classification of autism spectrum disorder using frequency-specific resting-state functional connectivity-A multi-center study. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 64, 1–9.
- Cima, M., Raine, A., Meesters, C., & Popma, A. (2013). Validation of the Dutch Reactive Proactive Questionnaire (RPQ): Differential Correlates of Reactive and Proactive Aggression From Childhood to Adulthood. *Aggressive Behavior*, 39(2), 99–113.
- Ciucci, E., Baroncelli, A., Golmaryami, F. N., & Frick, P. J. (2015). The Emotional Correlates to Callous-Unemotional Traits in Children. *Journal of Child and Family Studies*, 24(8), 2374–2387.
- Cognition, C. (1996). Cambridge neuropsychological test automated battery (CANTAB). CeNeS Limited, Cambridge.
- Cohn, M. D., Pape, L. E., Schmaal, L., van den Brink, W., van Wingen, G., Vermeiren, R. R. J. M., ... Popma, A. (2015). Differential relations between juvenile psychopathic traits and resting state network connectivity. *Human Brain Mapping*, 36(6), 2396–2405.
- Cohn, M. D., Viding, E., McCrory, E., Pape, L., van den Brink, W., Doreleijers, T. A. H., ... Popma, A. (2016). Regional grey matter volume and concentration in at-risk adolescents: Untangling associations with callous-unemotional traits and conduct disorder symptoms. *Psychiatry Research - Neuroimaging*, 254, 180–187.
- Connor, D. F., Steingard, R. J., Anderson, J. J., & Melloni, R. H. (2003). Gender differences in reactive and proactive aggression. *Child Psychiatry & Human Development*, 33(4), 279–294.
- Cools, R., Clark, L., Owen, A. M., & Robbins, T. W. (2002). Defining the neural mechanisms of probabilistic reversal learning using event-related functional magnetic resonance imaging. *The Journal of Neuroscience*, 22(11), 4563–4567.
- Crowley, T. J., Dalwani, M. S., Mikulich-Gilbertson, S. K., Du, Y. P., Lejuez, C. W., Raymond, K. M., & Banich, M. T. (2010). Risky decisions and their consequences: Neural processing by boys with antisocial substance disorder. *PLoS ONE*, 5(9), 1–20.
- Dadds, M. R., Cauchi, A. J., Wimalaweera, S., Hawes, D. J., & Brennan, J. (2012). Outcomes, moderators, and mediators of empathic-emotion recognition training for complex conduct problems in childhood. *Psychiatry Research*, 199(3), 201–207.
- Dadds, M. R., Fraser, J., Frost, A., & Hawes, D. J. (2005). Disentangling the

- underlying dimensions of psychopathy and conduct problems in childhood: a community study. *Journal of Consulting and Clinical Psychology*, 73(3), 400–410.
- Dadds, M. R., Kimonis, E. R., Schollar-Root, O., Moul, C., & Hawes, D. J. (2017). Are impairments in emotion recognition a core feature of callous-unemotional traits? Testing the primary versus secondary variants model in children. *Development and Psychopathology*, 1–11.
- Dalwani, M. S., Tregellas, J. R., Andrews-Hanna, J. R., Mikulich-Gilbertson, S. K., Raymond, K. M., Banich, M. T., ... Sakai, J. T. (2014). Default mode network activity in male adolescents with conduct and substance use disorder. *Drug and Alcohol Dependence*, 134(1), 242–250.
- Damoiseaux, J. S., Rombouts, S. A. R. B., Barkhof, F., Scheltens, P., Stam, C. J., Smith, S. M., & Beckmann, C. F. (2006). Consistent resting-state networks across healthy subjects. *Proceedings of the National Academy of Sciences*, 103(37), 13848–13853.
- Demir-Lira, Ö. E., Voss, J. L., O'Neil, J. T., Briggs-Gowan, M. J., Wakschlag, L. S., & Booth, J. R. (2016). Early-life stress exposure associated with altered prefrontal resting-state fMRI connectivity in young children. *Developmental Cognitive Neuroscience*, 19, 107–114.
- Deshpande, G., LaConte, S., Peltier, S., & Hu, X. (2009). Integrated local correlation: A new measure of local coherence in fMRI data. *Human Brain Mapping*, 30(1), 13–23.
- Diamond, A. (2013). Executive functions in attention-deficit/hyperactivity disorder. *Annual Review of Psychology*, 64, 135–168.
- Dipasquale, O., Sethi, A., Laganà, M. M., Baglio, F., Baselli, G., Kundu, P., ... Cercignani, M. (2017). Comparing resting state fMRI de-noising approaches using multi- and single-echo acquisitions. *Plos One*, 12(3), e0173289.
- Docherty, M., Boxer, P., Huesmann, L. R., O'Brien, M., & Bushman, B. (2017). Assessing Callous-Unemotional Traits in Adolescents: Determining Cutoff Scores for the Inventory of Callous and Unemotional Traits. *Journal of Clinical Psychology*, 73(3), 257–278.
- Dodge, K. a, & Coie, J. D. (1987). Social-information-processing factors in reactive and proactive aggression in children's peer groups. *Journal of Personality and Social Psychology*, 53(6), 1146–1158.
- Eisenbarth, H., Demetriou, C. A., Kyranides, M. N., & Fanti, K. A. (2016). Stability Subtypes of Callous-Unemotional Traits and Conduct Disorder Symptoms and Their Correlates. *Journal of Youth and*

- Adolescence*, 45(9), 1889–1901.
- Eisner, M., & Ribeaud, D. (2007). Conducting a criminological survey in a culturally diverse context: Lessons from the Zurich project on the social development of children. *European Journal of Criminology*, 4(3), 271–298.
- Emmert, K., Kopel, R., Sulzer, J., Brühl, A. B., Berman, B. D., Linden, D. E. J., ... Haller, S. (2016). Meta-analysis of real-time fMRI neurofeedback studies using individual participant data: How is brain regulation mediated? *NeuroImage*, 124, 806–812.
- Erskine, H. E., Ferrari, A. J., Polanczyk, G. V., Moffitt, T. E., Murray, C. J. L., Vos, T., ... Scott, J. G. (2014). The global burden of conduct disorder and attention-deficit/hyperactivity disorder in 2010. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 55(4), 328–336.
- Erskine, H. E., Norman, R. E., Ferrari, A. J., Chan, G. C. K., Copeland, W. E., Whiteford, H. A., & Scott, J. G. (2016). Long-Term Outcomes of Attention-Deficit/Hyperactivity Disorder and Conduct Disorder: A Systematic Review and Meta-Analysis. *Journal of the American Academy of Child & Adolescent Psychiatry*, 55(10), 841–850.
- Espinoza, F. A., Vergara, V. M., Reyes, D., Anderson, N. E., Harenski, C. L., Decety, J., ... Calhoun, V. D. (2018). Aberrant functional network connectivity in psychopathy from a large (N = 985) forensic sample. *Human Brain Mapping*, (February), 2624–2634.
- Essau, C. A., Sasagawa, S., & Frick, P. J. (2006). Callous-unemotional traits in a community sample of adolescents. *Assessment*, 13(4), 454–469.
- Fairchild, G., Hagan, C. C., Walsh, N. D., Passamonti, L., Calder, A. J., & Goodyer, I. M. (2013). Brain structure abnormalities in adolescent girls with conduct disorder. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 54(1), 86–95.
- Fairchild, G., Toschi, N., Hagan, C. C., Goodyer, I. M., Calder, A. J., & Passamonti, L. (2015). Cortical thickness, surface area, and folding alterations in male youths with conduct disorder and varying levels of callous-unemotional traits. *NeuroImage: Clinical*, 8, 253–260.
- Fanning, J. R., Keedy, S., Berman, M. E., Lee, R., & Coccaro, E. F. (2017). Neural Correlates of Aggressive Behavior in Real Time: a Review of fMRI Studies of Laboratory Reactive Aggression. *Current Behavioral Neuroscience Reports*, 4(2), 138–150.
- Fanti, K. A. (2016). Understanding heterogeneity in conduct disorder: A review of psychophysiological studies. *Neuroscience & Biobehavioral*

Reviews.

- Fanti, K. A., Demetriou, C. A., & Kimonis, E. R. (2013). Variants of Callous-Unemotional Conduct Problems in a Community Sample of Adolescents. *Journal of Youth and Adolescence*, 42(7), 964–979.
- Fanti, K. A., Panayiotou, G., Lazarou, C., Michael, R., & Georgiou, G. (2016). The better of two evils? Evidence that children exhibiting continuous conduct problems high or low on callous-unemotional traits score on opposite directions on physiological and behavioral measures of fear. *Development and Psychopathology*, 28(01), 185–198.
- Feilhauer, J., Cima, M., & Arntz, A. (2012). Assessing callous-unemotional traits across different groups of youths: Further cross-cultural validation of the Inventory of Callous-Unemotional Traits. *International Journal of Law and Psychiatry*, 35(4), 251–262.
- Finger, E. C., Marsh, A. A., Blair, K. S., Reid, M. E., Sims, C., Ng, P., ... Blair, R. J. R. (2011). Disrupted Reinforcement Signaling in Orbital Frontal Cortex and Caudate in Youths with Conduct Disorder/Oppositional Defiant Disorder and High Psychopathic Traits. *The American Journal of Psychiatry*, 168(2), 152–162.
- Fite, P. J., Colder, C. R., Lochman, J. E., & Wells, K. C. (2008). Developmental trajectories of proactive and reactive aggression from fifth to ninth grade. *Journal of Clinical Child and Adolescent Psychology*, 37(2), 412–421.
- Fite, P. J., Raine, A., Stouthamer-Loeber, M., Loeber, R., & Pardini, D. A. (2010). Reactive and proactive aggression in adolescent males: Examining differential outcomes 10 years later in early adulthood. *Criminal Justice and Behavior*, 37(2), 141–157.
- Fite, P. J., Rubens, S. L., Preddy, T. M., Raine, A., & Pardini, D. A. (2014). Reactive/proactive aggression and the development of internalizing problems in males: The moderating effect of parent and peer relationships. *Aggressive Behavior*, 40(1), 69–78.
- Fite, P. J., Stoppelbein, L., & Greening, L. (2009). Proactive and reactive aggression in a child psychiatric inpatient population: Relations to psychopathic characteristics. *Criminal Justice and Behavior*, 36(5), 481–493.
- Fonagy, P., & Luyten, P. (2017). Conduct problems in youth and the RDoC approach: A developmental, evolutionary-based view. *Clinical Psychology Review*.
- Fox, K. C. R., Spreng, R. N., Ellamil, M., Andrews-Hanna, J. R., & Christoff, K.

- (2015). The wandering brain: Meta-analysis of functional neuroimaging studies of mind-wandering and related spontaneous thought processes. *NeuroImage*, 111, 611–621.
- Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005). From The Cover: The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of Sciences*, 102(27), 9673–9678.
- Frick, P. J. (2004). *The Inventory of Callous-Unemotional Traits*. New Orleans: The University of New Orleans.
- Frick, P. J. (2012). Developmental pathways to conduct disorder: implications for future directions in research, assessment, and treatment. *Journal of Clinical Child and Adolescent Psychology*, 41(3), 378–389.
- Frick, P. J. (2016). Current research on conduct disorder in children and adolescents. *South African Journal of Psychology*, 46(2), 160–174.
- Frick, P. J., Ray, J. V., Thornton, L. C., & Kahn, R. E. (2013). Can callous-unemotional traits enhance the understanding, diagnosis, and treatment of serious conduct problems in children and adolescents? A comprehensive review. *Psychological Bulletin*, 140(1), 1–57.
- Frick, P. J., & White, S. F. (2008). Research Review: The importance of callous-unemotional traits for developmental models of aggressive and antisocial behavior. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 49(4), 359–375.
- Fumagalli, M., & Priori, A. (2012). Functional and clinical neuroanatomy of morality. *Brain*, 135(7), 2006–2021.
- Fung, A. L. C., Gerstein, L. H., Chan, Y., & Engebretson, J. (2015). Relationship of Aggression to Anxiety, Depression, Anger, and Empathy in Hong Kong. *Journal of Child and Family Studies*, 24(3), 821–831.
- Gao, Y., Tuvblad, C., Schell, A., Baker, L., & Raine, A. (2015). Skin conductance fear conditioning impairments and aggression: A longitudinal study. *Psychophysiology*, 52(2), 288–295.
- Goffin, K. C., Boldt, L. J., Kim, S., & Kochanska, G. (2017). A Unique Path to Callous-Unemotional Traits for Children who are Temperamentally Fearless and Unconcerned about Transgressions: a Longitudinal Study of Typically Developing Children from age 2 to 12. *Journal of Abnormal Child Psychology*, 1–12.
- Gomez, R., Vance, A., & Gomez, R. M. (2014). Analysis of the Convergent and Discriminant Validity of the CBCL, TRF, and YSR in a Clinic-Referred

- Sample. *Journal of Abnormal Child Psychology*, 42(8), 1413–1425.
- González-Madruga, K., Rogers, J., Toschi, N., Riccelli, R., Smaragdi, A., Puzzo, I., ... Fairchild, G. (2019). White matter microstructure of the extended limbic system in male and female youth with conduct disorder. *Psychological Medicine*, 1–10.
- Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38(5), 581–586.
- Greicius, M. D., Krasnow, B., Reiss, A. L., & Menon, V. (2003). Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proceedings of the National Academy of Sciences of the United States of America*, 100(1), 253–258.
- Griffiths, S. Y., & Jalava, J. V. (2017). A comprehensive neuroimaging review of PCL-R defined psychopathy. *Aggression and Violent Behavior*.
- Guelker, M. D., Barry, C. T., Barry, T. D., & Malkin, M. L. (2014). Perceived positive outcomes as a mediator between adolescent callous-unemotional traits and antisocial behavior. *Personality and Individual Differences*, 69, 129–134.
- Haney-Caron, E., Caprihan, A., & Stevens, M. C. (2014). DTI-measured white matter abnormalities in adolescents with Conduct Disorder. *Journal of Psychiatric Research*, 48(1), 111–120.
- Hartley, C. M., Pettit, J. W., & Castellanos, D. (2016). Reactive Aggression and Suicide-Related Behaviors in Children and Adolescents: A Review and Preliminary Meta-Analysis. *Suicide and Life-Threatening Behavior*, 1–14.
- Herpers, P. C. M., Scheepers, F. E., Bons, D. M. a, Buitelaar, J. K., & Rommelse, N. N. J. (2014). The cognitive and neural correlates of psychopathy and especially callous-unemotional traits in youths: a systematic review of the evidence. *Development and Psychopathology*, 26(1), 245–273.
- Holz, N. E., Boecker-Schlier, R., Buchmann, A. F., Blomeyer, D., Jennen-Steinmetz, C., Baumeister, S., ... Laucht, M. (2016). Ventral striatum and amygdala activity as convergence sites for early adversity and conduct disorder. *Social Cognitive and Affective Neuroscience*, (September 2016), nsw120.
- Huettel, S. a., Song, A. W., & McCarthy, G. (2009). *Functional Magnetic Resonance Imaging*.
- Humayun, S., Kahn, R. E., Frick, P. J., & Viding, E. (2014). Callous-Unemotional Traits and Anxiety in a Community Sample of 7-Year-Olds. *Journal of Clinical Child and Adolescent Psychology*, 43(1), 36–42.

- Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., ... Wang, P. (2010). Research Domain Criteria (RDoC): Toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*, 167(7), 748–751.
- Jambon, M., & Smetana, J. G. (2018). Individual Differences in Prototypical Moral and Conventional Judgments and Children's Proactive and Reactive Aggression. *Child Development*, 89(4), 1343–1359.
- Jovicich, J., Minati, L., Marizzoni, M., Marchitelli, R., Sala-Llonch, R., Bartrés-Faz, D., ... Frisoni, G. B. (2016). Longitudinal reproducibility of default-mode network connectivity in healthy elderly participants: A multicentric resting-state fMRI study. *NeuroImage*, 124, 442–454.
- Kahn, R. E., Ermer, E., Salovey, P., & Kiehl, K. A. (2016). Emotional Intelligence and Callous-Unemotional Traits in Incarcerated Adolescents. *Child Psychiatry and Human Development*, 47(6), 903–917.
- Kahn, R. E., Frick, P. J., Golmaryami, F. N., & Marsee, M. A. (2017). The Moderating Role of Anxiety in the Associations of Callous-Unemotional Traits with Self-Report and Laboratory Measures of Affective and Cognitive Empathy. *Journal of Abnormal Child Psychology*, 45(3), 583–596.
- Kaufman, J., Birmaher, B., Brent, D., Rao, U., Flynn, C., Moreci, P., ... Ryan, N. (1997). Schedule for affective disorders and schizophrenia for school-age children-present and lifetime version (K-SADS-PL): Initial reliability and validity data. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(7), 980–988.
- Kay, S. R., Wolkenfeld, F., & Murrill, L. M. (1988). Profiles of aggression among psychiatric patients. *The Journal of Nervous and Mental Disease*, 176(9), 539–546.
- Kiehl, K. A. (2006). A cognitive neuroscience perspective on psychopathy: Evidence for paralimbic system dysfunction. *Psychiatry Research*, 142(2–3), 107–128.
- Kimonis, E. R., Frick, P. J., Cauffman, E., Goldweber, A., & Skeem, J. L. (2012). Primary and secondary variants of juvenile psychopathy differ in emotional processing. *Development and Psychopathology*, 24(3), 1091–1103.
- Kimonis, E. R., Frick, P. J., Skeem, J. L., Marsee, M. A., Cruise, K., Centifanti, L. C., ... Morris, A. S. (2008). Assessing callous-unemotional traits in adolescent offenders: Validation of the Inventory of Callous-

- Unemotional Traits. *International Journal of Law and Psychiatry*, 31(3), 241–252.
- Korponay, C., Pujara, M., Deming, P., Philippi, C., Decety, J., Kosson, D. S., ... Koenigs, M. (2017). Impulsive-Antisocial Dimension of Psychopathy Linked to Enlargement and Abnormal Functional Connectivity of the Striatum. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 2(2), 149–157.
- Kundu, P., Voon, V., Balchandani, P., Lombardo, M. V., Poser, B. A., & Bandettini, P. A. (2017). Multi-echo fMRI: A review of applications in fMRI denoising and analysis of BOLD signals. *NeuroImage*, 154(March), 59–80.
- Latzman, R. D., Lilienfeld, S. O., Latzman, N. E., & Clark, L. A. (2013). Exploring callous and unemotional traits in youth via general personality traits: An eye toward DSM-5. *Personality Disorders: Theory, Research, and Treatment*, 4(3), 191–202.
- Lee, M. H., Smyser, C. D., & Shimony, J. S. (2013). Resting state fMRI: A review of methods and clinical applications. *AJNR Am J Neuroradiol*, 34(10), 1866–1872.
- Leung, P. W. L., Kwong, S. L., Tang, C. P., Ho, T. P., Hung, S. F., Lee, C. C., ... Liu, W. S. (2006). Test-retest reliability and criterion validity of the Chinese version of CBCL, TRF, and YSR. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 47(9), 970–973.
- Lilienfeld, S. O. (2014). The Research Domain Criteria (RDoC): An analysis of methodological and conceptual challenges. *Behaviour Research and Therapy*, 62, 129–139.
- Lindquist, K. A., Wager, T. D., Kober, H., Bliss-Moreau, E., & Barrett, L. F. (2012). The brain basis of emotion: A meta-analytic review. *Behavioral and Brain Sciences*, 35(3), 121–143.
- Liu, H., Stufflebeam, S. M., Sepulcre, J., Hedden, T., & Buckner, R. L. (2009). Evidence from intrinsic activity that asymmetry of the human brain is controlled by multiple factors. *Proceedings of the National Academy of Sciences of the United States of America*, 106(48), 20499–20503.
- Lockwood, P. L., Sebastian, C. L., McCrory, E. J., Hyde, Z. H., Gu, X., De Brito, S. A., & Viding, E. (2013). Association of callous traits with reduced neural response to others' pain in children with conduct problems. *Current Biology*, 23(10), 901–905.
- Loeber, R., Burke, J. D., Lahey, B. B., Winters, A., & Zera, M. (2000). Oppositional Defiant and Conduct Disorder: A Review of the Past 10

- Years, Part I. *Journal of the American Academy of Child & Adolescent Psychiatry*, 39(12), 1468–1484.
- Loeber, R., Burke, J., & Pardini, D. A. (2009). Perspectives on oppositional defiant disorder, conduct disorder, and psychopathic features. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 50(1–2), 133–142.
- Longman, T., Hawes, D. J., & Kohlhoff, J. (2016). Callous–Unemotional Traits as Markers for Conduct Problem Severity in Early Childhood: A Meta-analysis. *Child Psychiatry and Human Development*, 47(2), 326–334.
- Lozier, L. M., Cardinale, E. M., VanMeter, J. W., & Marsh, A. A. (2014). Mediation of the Relationship Between Callous-Unemotional Traits and Proactive Aggression by Amygdala Response to Fear Among Children With Conduct Problems. *JAMA Psychiatry*, 71(6), 627.
- Lu, F.-M., Zhou, J.-S., Wang, X.-P., Xiang, Y.-T., & Yuan, Z. (2017). Short- and long-range functional connectivity density alterations in adolescents with pure conduct disorder at resting-state. *Neuroscience*, 351, 96–107.
- Lu, F.-M., Zhou, J.-S., Zhang, J., Wang, X., & Yuan, Z. (2017). Disrupted small-world brain network topology in pure conduct disorder. *Oncotarget, Advance Pu*.
- Lu, F.-M., Zhou, J.-S., Zhang, J., Xiang, Y.-T., Zhang, J., Liu, Q., ... Yuan, Z. (2015). Functional Connectivity Estimated from Resting-State fMRI Reveals Selective Alterations in Male Adolescents with Pure Conduct Disorder. *PLoS ONE*, 10(12), e0145668.
- Mak, L. E., Minuzzi, L., MacQueen, G., Hall, G., Kennedy, S. H., & Milev, R. (2017). The Default Mode Network in Healthy Individuals: A Systematic Review and Meta-Analysis. *Brain Connectivity*, 7(1), 25–33.
- Marsee, M. A., Barry, C. T., Childs, K. K., Frick, P. J., Kimonis, E. R., Centifanti, L. C., ... Lau, K. S. (2011). Assessing the forms and functions of aggression using self-report: factor structure and invariance of the peer conflict scale in youths. *Psychological Assessment*, 23(3), 792–804.
- Marsee, M. A., & Frick, P. J. (2007). Exploring the Cognitive and Emotional Correlates to Proactive and Reactive Aggression in a Sample of Detained Girls. *Journal of Abnormal Child Psychology*, 35(6), 969–981.
- Marsee, M. A., Weems, C. F., & Taylor, L. K. (2008). Exploring the association between aggression and anxiety in youth: A look at aggressive subtypes, gender, and social cognition. *Journal of Child and Family*

- Studies*, 17(1), 154–168.
- Marsh, A. a, Finger, E. C., Mitchell, D. G. V, Reid, M. E., Sims, C., Kosson, D. S., ... Blair, R. J. R. (2008). Reduced amygdala response to fearful expressions in children and adolescents with callous-unemotional traits and disruptive behavior disorders.[Erratum appears in *Am J Psychiatry*. 2008 Jul;165(7):920], [Erratum appears in *Am J Psychiatry*. 2008 May;165(5): *American Journal of Psychiatry*, 165(6), 712–720.
- Martuzzi, R., Ramani, R., Qiu, M., Shen, X., Papademetris, X., & Constable, R. T. (2011). A whole-brain voxel based measure of intrinsic connectivity contrast reveals local changes in tissue connectivity with anesthetic without a priori assumptions on thresholds or regions of interest. *NeuroImage*, 58(4), 1044–1050.
- Matthys, W., Vanderschuren, L. J. M. J., & Schutter, D. J. L. G. (2013). The neurobiology of oppositional defiant disorder and conduct disorder: Altered functioning in three mental domains. *Development and Psychopathology*, 25, 1–15.
- McMahon, R. J., Witkiewitz, K., & Kotler, J. S. (2010). Predictive validity of callous-unemotional traits measured in early adolescence with respect to multiple antisocial outcomes. *Journal of Abnormal Psychology*, 119(4), 752–763.
- Meyer-Lindenberg, A. (2010). From maps to mechanisms through neuroimaging of schizophrenia. *Nature*, 468(7321), 194–202.
- Michalska, K. J., Zeffiro, T. A., & Decety, J. (2016). Brain response to viewing others being harmed in children with conduct disorder symptoms. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 57(4), 510–519.
- Moffitt, T. E. (1993). Adolescence-limited and life-course-persistent antisocial behavior: a developmental taxonomy. *Psychological Review*, 100(4), 674–701.
- Moffitt, T. E., Caspi, A., Harrington, H., & Milne, B. J. (2002). Males on the life-course-persistent and adolescence-limited antisocial pathways: follow-up at age 26 years. *Development and Psychopathology*, 14(1), 179–207.
- Motzkin, J. C., Newman, J. P., Kiehl, K. A., & Koenigs, M. (2011). Reduced prefrontal connectivity in psychopathy. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 31(48), 17348–17357.

- Moul, C., Hawes, D. J., & Dadds, M. R. (2016). Mapping the developmental pathways of child conduct problems through the neurobiology of empathy. *Neuroscience and Biobehavioral Reviews*, 91, 34–50.
- Muehlhan, M., Lueken, U., Wittchen, H. U., & Kirschbaum, C. (2011). The scanner as a stressor: Evidence from subjective and neuroendocrine stress parameters in the time course of a functional magnetic resonance imaging session. *International Journal of Psychophysiology*, 79(2), 118–126.
- Muñoz, L. C. (2009). Callous-unemotional traits are related to combined deficits in recognizing afraid faces and body poses. *Journal of the American Academy of Child and Adolescent Psychiatry*, 48(5), 554–562.
- Muratori, P., Lochman, J. E., Manfredi, A., Milone, A., Nocentini, A., Pisano, S., & Masi, G. (2016). Callous unemotional traits in children with disruptive behavior disorder: Predictors of developmental trajectories and adolescent outcomes. *Psychiatry Research*, 236, 35–41.
- Nagai, Y., Critchley, H. D., Featherstone, E., Trimble, M. R., & Dolan, R. J. (2004). Activity in ventromedial prefrontal cortex covaries with sympathetic skin conductance level: A physiological account of a “default mode” of brain function. *NeuroImage*, 22(1), 243–251.
- Noordermeer, S. D. S., Luman, M., & Oosterlaan, J. (2016). A Systematic Review and Meta-analysis of Neuroimaging in Oppositional Defiant Disorder (ODD) and Conduct Disorder (CD) Taking Attention-Deficit Hyperactivity Disorder (ADHD) Into Account. *Neuropsychology Review*, 1–29.
- Ogawa, S., Lee, T. M., Kay, A. R., & Tank, D. W. (1990). Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proceedings of the National Academy of Sciences of the United States of America*, 87(24), 9868–9872.
- Ormel, J., Oldehinkel, A. J., Sijtsma, J., van Oort, F., Raven, D., Veenstra, R., ... Verhulst, F. C. (2012). The Tracking Adolescents' Individual Lives Survey (TRAILS): Design, current status, and selected findings. *Journal of the American Academy of Child and Adolescent Psychiatry*, 51(10), 1020–1036.
- Pardini, D. A., Raine, A., Erickson, K., & Loeber, R. (2014). Lower amygdala volume in men is associated with childhood aggression, early psychopathic traits, and future violence. *Biological Psychiatry*, 75(1), 73–80.

- Pechorro, P., Ray, J. V., Gonçalves, R. A., & Jesus, S. N. (2017). The Inventory of Callous-Unemotional Traits: Psychometric properties among referred and non-referred Portuguese female juveniles. *International Journal of Law and Psychiatry*, 54, 67–75.
- Peraza, J., Cservénka, A., Herting, M. M., & Nagel, B. J. (2015). Atypical parietal lobe activity to subliminal faces in youth with a family history of alcoholism. *American Journal of Drug and Alcohol Abuse*, 41(2), 139–145.
- Perenc, L., & Radochonski, M. (2014). Psychopathic traits and reactive-proactive aggression in a large community sample of Polish adolescents. *Child Psychiatry and Human Development*, 45(4), 464–471.
- Pessoa, L., Kastner, S., & Ungerleider, L. G. (2002). Attentional control of the processing of neural and emotional stimuli. *Brain Research. Cognitive Brain Research*, 15(1), 31–45.
- Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*, 17(2), 117–133.
- Philippi, C. L., Pujara, M. S., Motzkin, J. C., Newman, J., Kiehl, K. A., & Koenigs, M. (2015). Altered Resting-State Functional Connectivity in Cortical Networks in Psychopathy. *Journal of Neuroscience*, 35(15), 6068–6078.
- Pihet, S., Etter, S., Schmid, M., & Kimonis, E. R. (2015). Assessing Callous-Unemotional Traits in Adolescents: Validity of the Inventory of Callous-Unemotional Traits Across Gender, Age, and Community/Institutionalized Status. *Journal of Psychopathology and Behavioral Assessment*, 37(3), 407–421.
- Polanczyk, G. V., Salum, G. A., Sugaya, L. S., Caye, A., & Rohde, L. A. (2015). Annual research review: A meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *Journal of Child Psychology and Psychiatry*, 56(3), 345–365.
- Polman, H., Orobio De Castro, B., Kooops, W., Van Boxtel, H. W., & Merk, W. W. (2007). A meta-analysis of the distinction between reactive and proactive aggression in children and adolescents. *Journal of Abnormal Child Psychology*, 35(4), 522–535.
- Porter, D. (2018). RDoC, Psychopathology, and Naturalism: What's New Is What's Old? *Journal of Humanistic Psychology*.
- Poulin, F., & Boivin, M. (2000). Reactive and proactive aggression: evidence

- of a two factor-model. *Psychological Assessment*, 12(2), 115–122.
- Pruim, R. H. R., Mennes, M., Buitelaar, J. K., & Beckmann, C. F. (2015). Evaluation of ICA-AROMA and alternative strategies for motion artifact removal in resting state fMRI. *NeuroImage*, 112, 278–287.
- Pu, W., Luo, Q., Jiang, Y., Gao, Y., Ming, Q., & Yao, S. (2017). Alterations of Brain Functional Architecture Associated with Psychopathic Traits in Male Adolescents with Conduct Disorder. *Scientific Reports*, 7(1), 11349.
- Puiu, A. A., Wudarczyk, O., Goerlich, K. S., Votinov, M., Herpertz-dahlmann, B., Turetsky, B., & Konrad, K. (2018). Neuroscience and Biobehavioral Reviews Impulsive aggression and response inhibition in attention-deficit / hyperactivity disorder and disruptive behavioral disorders : Findings from a systematic review. *Neuroscience and Biobehavioral Reviews*, 90(March), 231–246.
- Pujol, J., Batalla, I., Contreras-Rodríguez, O., Harrison, B. J., Pera, V., Hernández-Ribas, R., ... Cardoner, N. (2012). Breakdown in the brain network subserving moral judgment in criminal psychopathy. *Social Cognitive and Affective Neuroscience*, 7(8), 917–923.
- Puzzo, I., Smaragdi, A., Gonzalez, K., Martin-Key, N., & Fairchild, G. (2016). Neurobiological, Neuroimaging, and Neuropsychological Studies of Children and Adolescents with Disruptive Behavior Disorders. *Family Relations*, 65, 134–150.
- Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001). A default mode of brain function. *Proceedings of the National Academy of Sciences of the United States of America*, 98(2), 676–682.
- Raine, A. (2018). Antisocial Personality as a Neurodevelopmental Disorder. *Annual Review of Clinical Psychology*, 14(1), annurev-clinpsy-050817-084819.
- Raine, A., Dodge, K., Loeber, R., Gatzke-Kopp, L., Lynam, D., Reynolds, C., ... Liu, J. (2006). The Reactive–Proactive Aggression Questionnaire: Differential Correlates of Reactive and Proactive Aggression in Adolescent Boys. *Aggressive Behaviour*, 32(3), 159–171.
- Raine, A., Dodge, K., Loeber, R., Gatzke-Kopp, L., Lynam, D., Stouthamer-Loeber, M., & Liu, J. (2006). The Reactive–Proactive Aggression Questionnaire: Differential Correlates of Reactive and Proactive Aggression in Adolescent Boys. *Aggressive Behav.*, 32(2), 159–171.
- Raine, A., Reynolds, C., Venables, P. H., Mednick, S. A., & Farrington, D. P.

- (1998). Fearlessness, stimulation-seeking, and large body size at age 3 years as early predispositions to childhood aggression at age 11 years. *Archives of General Psychiatry*, 55(8), 745–751.
- Raine, A., & Yang, Y. (2006). Neural foundations to moral reasoning and antisocial behavior. *Social Cognitive and Affective Neuroscience*, 1(3), 203–213.
- Raschle, N. M., Menks, W. M., Fehlbauer, L. V., Tshomba, E., & Stadler, C. (2015). Structural and functional alterations in right dorsomedial prefrontal and left insular cortex co-localize in adolescents with aggressive behaviour: An ALE meta-analysis. *PLoS ONE*, 10(9), 1–24.
- Reidy, D. E., Krusemark, E., Kosson, D. S., Kearns, M. C., Smith-Darden, J., & Kiehl, K. A. (2017). The Development of Severe and Chronic Violence Among Youth: The Role of Psychopathic Traits and Reward Processing. *Child Psychiatry and Human Development*, 48(6), 967–982.
- Ridderinkhof, K. R., Ullsperger, M., Crone, E. A., & Nieuwenhuis, S. (2004). The Role of the Medial Frontal Cortex in Cognitive Control. *Science*, 306(5695), 443–447.
- Rogers, J. C., & De Brito, S. A. (2016). Cortical and Subcortical Gray Matter Volume in Youths With Conduct Problems. *JAMA Psychiatry*, 73(1), 64.
- Rosazza, C., Minati, L., Ghielmetti, F., Mandelli, M. L., & Bruzzone, M. G. (2012). Functional Connectivity during Resting-State Functional MR Imaging: Study of the Correspondence between Independent Component Analysis and Region-of-Interest-Based Methods. *AJNR Am J Neuroradiol*, 33, 180–187.
- Roy, A. K., Bennett, R., Posner, J., Hulvershorn, L., Castellanos, F. X., & Klein, R. G. (2018). Altered intrinsic functional connectivity of the cingulate cortex in children with severe temper outbursts. *Development and Psychopathology*, 30(2), 571–579.
- Rubia, K., Halari, R., Mohammad, A. M., Taylor, E., & Brammer, M. (2011). Methylphenidate normalizes frontocingulate underactivation during error processing in attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 70(3), 255–262.
- Saad, Z. S., Reynolds, R. C., Jo, H. J., Gotts, S. J., Chen, G., Martin, A., & Cox, R. W. (2013). Correcting Brain-Wide Correlation Differences in Resting-State FMRI. *Brain Connectivity*, 3(4), 339–352.
- Salihovic, S., & Stattin, H. (2017). Psychopathic Traits and Delinquency Trajectories in Adolescence. *Journal of Psychopathology and Behavioral Assessment*, 39(1), 15–24.

- Salvatore, J. E., & Dick, D. M. (2016). Genetic influences on conduct disorder. *Neuroscience & Biobehavioral Reviews*.
- Sato, J. R., Biazoli, C. E., Salum, G. A., Gadelha, A., Crossley, N., Vieira, G., ... Bressan, R. A. (2018). Association between abnormal brain functional connectivity in children and psychopathology: A study based on graph theory and machine learning. *World Journal of Biological Psychiatry*, 19(2), 119–129.
- Scarpa, A., Haden, S. C., & Tanaka, A. (2010). Being hot-tempered: Autonomic, emotional, and behavioral distinctions between childhood reactive and proactive aggression. *Biological Psychology*, 84(3), 488–496.
- Schoenberg, P. L. A., & David, A. S. (2014). Biofeedback for psychiatric disorders: a systematic review. *Applied Psychophysiology and Biofeedback*, 39(2), 109–135.
- Sebastian, C. L., McCrory, E. J., Dadds, M. R., Cecil, C. A. M., Lockwood, P. L., Hyde, Z. H., ... Viding, E. (2014). Neural responses to fearful eyes in children with conduct problems and varying levels of callous-unemotional traits. *Psychological Medicine*, 44(01), 99–109.
- Sethi, A., McCrory, E., Puetz, V., Hoffmann, F., Knodt, A. R., Radtke, S. R., ... Viding, E. (2018). Primary and Secondary Variants of Psychopathy in a Volunteer Sample Are Associated With Different Neurocognitive Mechanisms. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 3(12), 1013–1021.
- Sethi, A., Sarkar, S., Dell’Acqua, F., Viding, E., Catani, M., Murphy, D. G. M., & Craig, M. C. (2018). Anatomy of the dorsal default-mode network in conduct disorder: Association with callous-unemotional traits. *Developmental Cognitive Neuroscience*, 30(September 2017), 87–92.
- Shannon, B. J., Raichle, M. E., Snyder, A. Z., Fair, D. a, Mills, K. L., Zhang, D., ... Kiehl, K. a. (2011). Premotor functional connectivity predicts impulsivity in juvenile offenders. *Proceedings of the National Academy of Sciences of the United States of America*, 108(27), 11241–11245.
- Sherman, L. E., Rudie, J. D., Pfeifer, J. H., Masten, C. L., Mcnealy, K., & Dapretto, M. (2014). Developmental Cognitive Neuroscience Development of the Default Mode and Central Executive Networks across early adolescence : A longitudinal study. *Accident Analysis and Prevention*, 10, 148–159.
- Siep, N., Tonnaer, F., van de Ven, V., Arntz, A., Raine, A., & Cima, M. (2018). Anger provocation increases limbic and decreases medial prefrontal

- cortex connectivity with the left amygdala in reactive aggressive violent offenders. *Brain Imaging and Behavior*.
- Silbersweig, D., & Loscalzo, J. (2017). Precision psychiatry meets network medicine network psychiatry. *JAMA Psychiatry*, 74(7), 665–666.
- Sitaram, R., Caria, A., Veit, R., Gaber, T., Ruiz, S., & Birbaumer, N. (2014). Volitional control of the anterior insula in criminal psychopaths using real-time fMRI neurofeedback: a pilot study. *Frontiers in Behavioral Neuroscience*, 8(October), 344.
- Slot, W., Orobio De Castro, B., & Duivenvoorden, Y. (2000). *WAS-lijst, eerste afname, versie 1*.
- Smaragdi, A., Cornwell, H., Toschi, N., Riccelli, R., Gonzalez-Madruga, K., Wells, A., ... Fairchild, G. (2017). Sex Differences in the Relationship Between Conduct Disorder and Cortical Structure in Adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 56(8), 703–712.
- Smeets, K. C., Oostermeijer, S., Lappenschaar, M., Cohn, M., van der Meer, J. M. J., Popma, A., ... Buitelaar, J. K. (2016). Are Proactive and Reactive Aggression Meaningful Distinctions in Adolescents? A Variable- and Person-Based Approach. *Journal of Abnormal Child Psychology*, 1–14.
- Smitha, K. A., Akhil Raja, K., Arun, K. M., Rajesh, P. G., Thomas, B., Kapilamoorthy, T. R., & Kesavadas, C. (2017). Resting state fMRI: A review on methods in resting state connectivity analysis and resting state networks. *Neuroradiology Journal*, 30(4), 305–317.
- Swanson, J. M. (1992). *School-based assessments and interventions for ADD students*. KC publishing.
- Sylvers, P. D., Brennan, P. a, & Lilienfeld, S. O. (2011). Psychopathic traits and preattentive threat processing in children: a novel test of the fearlessness hypothesis. *Psychological Science*, 22(10), 1280–1287.
- Szabó, E., Kocsel, N., Édes, A., Pap, D., Galambos, A., Zsombók, T., ... Kökönyei, G. (2017). Callous-unemotional traits and neural responses to emotional faces in a community sample of young adults. *Personality and Individual Differences*, 111, 312–317.
- Tang, Y., Jiang, W., Liao, J., Wang, W., & Luo, A. (2013). Identifying Individuals with Antisocial Personality Disorder Using Resting-State fMRI. *PLoS ONE*, 8(4).
- Tang, Y., Liu, W., Chen, J., Liao, J., Hu, D., & Wang, W. (2013). Altered spontaneous activity in antisocial personality disorder revealed by regional homogeneity. *NeuroReport*, 24(11), 590–595.

- Tang, Y., Long, J., Wang, W., Liao, J., Xie, H., Zhao, G., & Zhang, H. (2016). Aberrant functional brain connectome in people with antisocial personality disorder. *Scientific Reports*, 6(1), 26209.
- Thijssen, S., & Kiehl, K. A. (2017). Functional connectivity in incarcerated male adolescents with psychopathic traits. *Psychiatry Research - Neuroimaging*, 265, 35–44.
- Thomason, M. E., Dennis, E. L., Joshi, A. A., Joshi, S. H., Dinov, I. D., Chang, C., ... Gotlib, I. H. (2011). Resting-state fMRI can reliably map neural networks in children. *NeuroImage*, 55(1), 165–175.
- Tomasi, D., & Volkow, N. D. (2012). Resting functional connectivity of language networks: Characterization and reproducibility. *Molecular Psychiatry*, 17(8), 841–854.
- Tremblay, R. E., Vitaro, F., & Côté, S. M. (2018). Developmental Origins of Chronic Physical Aggression: A Bio-Psycho-Social Model for the Next Generation of Preventive Interventions. *Annual Review of Psychology*, 69(1), 383–407.
- Umbach, R., Berryessa, C. M., & Raine, A. (2015). Brain imaging research on psychopathy: Implications for punishment, prediction, and treatment in youth and adults. *Journal of Criminal Justice*, 43(4), 295–306.
- Urban, S., Habersaat, S., Pihet, S., Suter, M., de Ridder, J., & Stéphan, P. (2018). Specific Contributions of Age of Onset, Callous-Unemotional Traits and Impulsivity to Reactive and Proactive Aggression in Youths with Conduct Disorders. *Psychiatric Quarterly*, 89(1), 1–10.
- Uytun, M. C., Karakaya, E., Oztop, D. B., Gengec, S., Gumus, K., Ozmen, S., ... Ozsoy, S. D. (2016). Default mode network activity and neuropsychological profile in male children and adolescents with attention deficit hyperactivity disorder and conduct disorder. *Brain Imaging and Behavior*.
- van Dijk, K. R. A., Hedden, T., Venkataraman, A., Karleyton, C., Lazar, S. W., Buckner, R. L., ... Buckner, R. L. (2010). Intrinsic Functional Connectivity As a Tool For Human Connectomics : Theory , Properties , and Optimization Intrinsic Functional Connectivity As a Tool For Human Connectomics : Theory , Properties , and Optimization, 02138, 297–321.
- van Goozen, S. H. M., Fairchild, G., Snoek, H., & Harold, G. T. (2007). The evidence for a neurobiological model of childhood antisocial behavior. *Psychological Bulletin*, 133(1), 149–182.
- Van Ombergen, A., Heine, L., Jillings, S., Roberts, R. E., Jeurissen, B., Van

- Rompaey, V., ... Wuyts, F. L. (2017). Altered functional brain connectivity in patients with visually induced dizziness. *NeuroImage: Clinical*, 14, 538–545.
- Vatansever, D., Manktelow, A. E., Sahakian, B. J., Menon, D. K., & Stamatakis, E. A. (2017). Angular default mode network connectivity across working memory load. *Human Brain Mapping*, 38(1), 41–52.
- Veroude, K., von Rhein, D., Chauvin, R. J. M., van Dongen, E. V., Mennes, M. J. J., Franke, B., ... Buitelaar, J. K. (2016). The link between callous-unemotional traits and neural mechanisms of reward processing: An fMRI study. *Psychiatry Research - Neuroimaging*, 255, 75–80.
- Via, E., Cardoner, N., Pujol, J., Alonso, P., Lopez-Sola, M., Real, E., ... Harrison, B. J. (2014). Amygdala activation and symptom dimensions in obsessive-compulsive disorder. *The British Journal of Psychiatry*, 204(1), 61–68.
- Vitaro, F., Barker, E. D., Boivin, M., Brendgen, M., & Tremblay, R. E. (2006). Do early difficult temperament and harsh parenting differentially predict reactive and proactive aggression? *Journal of Abnormal Child Psychology*, 34(5), 685–695.
- Vitaro, F., Brendgen, M., & Tremblay, R. E. (2002). Reactively and proactively aggressive children: Antecedent and subsequent characteristics. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 43(4), 495–505.
- Vitiello, B., & Stoff, D. M. (1997). Subtypes of Aggression and Their Relevance to Child Psychiatry. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36(3), 307–315.
- Von Rhein, D., Cools, R., Zwiers, M. P., Van Der Schaaf, M., Franke, B., Luman, M., ... Buitelaar, J. (2015). Increased neural responses to reward in adolescents and young adults with attention-deficit/hyperactivity disorder and their unaffected siblings. *Journal of the American Academy of Child and Adolescent Psychiatry*, 54(5), 394–402.
- von Rhein, D., Oldehinkel, M., Beckmann, C. F., Oosterlaan, J., Heslenfeld, D., Hartman, C. A., ... Mennes, M. (2016). Aberrant local striatal functional connectivity in attention-deficit/hyperactivity disorder. *Journal of Child Psychology and Psychiatry*, 3, 1–9.
- Wallace, G. L., White, S. F., Robustelli, B., Sinclair, S., Hwang, S., Martin, A., & Blair, R. J. R. (2014). Cortical and subcortical abnormalities in youths with conduct disorder and elevated callous-unemotional traits. *Journal of the American Academy of Child and Adolescent Psychiatry*,

- 53(4), 456–465.e1.
- Waller, R., & Hyde, L. W. (2017). Callous–Unemotional Behaviors in Early Childhood: Measurement, Meaning, and the Influence of Parenting. *Child Development Perspectives*, 11(2), 120–126.
- Waller, R., & Hyde, L. W. (2018). Callous-unemotional behaviors in early childhood: the development of empathy and prosociality gone awry. *Current Opinion in Psychology*, 20, 11–16.
- Walpolá, I. C., Nest, T., Roseman, L., Erritzoe, D., Feilding, A., Nutt, D. J., & Carhart-Harris, R. L. (2017). Altered Insula Connectivity under MDMA. *Neuropsychopharmacology*, 42(11), 2152–2162.
- Waltes, R., Chiochetti, A. G., & Freitag, C. M. (2016). The neurobiological basis of human aggression: A review on genetic and epigenetic mechanisms. *American Journal of Medical Genetics, Part B: Neuropsychiatric Genetics*, 171(5), 650–675.
- Waschbusch, D. A. (2002). A meta-analytic examination of comorbid hyperactive-impulsive-attention problems and conduct problems. *Psychological Bulletin*, 128(1), 118–150.
- Wechsler, D. (2003). *Wechsler intelligence scale for children-WISC-IV*. Psychological Corporation.
- Werhahn, J. E., Mohl, S., Willinger, D., Smigielski, L., Roth, A., Mulder, L. M., ... Castro, J. (2018). Aggression Subtypes Relate to Distinct Resting State Functional Connectivity in Children and Adolescents with Disruptive Behavior, 1–25.
- Wesselhoeft, R., Stringaris, A., Sibbersen, C., Kristensen, R. V., Bojesen, A. B., & Talati, A. (2018). Dimensions and subtypes of oppositionality and their relation to comorbidity and psychosocial characteristics. *European Child and Adolescent Psychiatry*, (0123456789), 1–15.
- White, B. A., Gordon, H., & Guerra, R. C. (2015). Callous-unemotional traits and empathy in proactive and reactive relational aggression in young women. *Personality and Individual Differences*, 75, 185?189.
- Whitfield-Gabrieli, S., & Ford, J. M. (2012). Default Mode Network Activity and Connectivity in Psychopathology. *Annual Review of Clinical Psychology*, 8(1), 49–76.
- Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A Functional Connectivity Toolbox for Correlated and Anticorrelated Brain Networks. *Brain Connectivity*, 2(3), 125–141.
- Wilkinson, S., Waller, R., & Viding, E. (2016). Practitioner Review: Involving

- young people with callous unemotional traits in treatment - Does it work? A systematic review. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 57(5), 552–565.
- Wong, T. Y., Sid, A., Wensing, · Tobias, Eickhoff, S. B., Habel, U., Gur, R. C., & Nickl-Jockschat, T. (2018). Neural networks of aggression: ALE meta-analyses on trait and elicited aggression. *Brain Structure and Function*, 0(0), 0.
- Woodward, N. D., & Cascio, C. J. (2015). Resting-State Functional Connectivity in Psychiatric Disorders. *JAMA Psychiatry*, 72(8), 743–744.
- Wu, Q., Zhang, X., Dong, D., Wang, X., & Yao, S. (2017). Altered spontaneous brain activity in adolescent boys with pure conduct disorder revealed by regional homogeneity analysis. *European Child and Adolescent Psychiatry*, 26(7), 1–11.
- Yang, Y., Narr, K. L., Baker, L. A., Joshi, S. H., Jahanshad, N., Raine, A., & Thompson, P. M. (2015). Frontal and striatal alterations associated with psychopathic traits in adolescents. *Psychiatry Research - Neuroimaging*, 231(3), 333–340.
- Yendiki, A., Greve, D. N., Wallace, S., Vangel, M., Bockholt, J., Mueller, B. A., ... Gollub, R. L. (2010). Multi-site characterization of an fMRI working memory paradigm: Reliability of activation indices. *NeuroImage*, 53(1), 119–131.
- Yoder, K. J., Lahey, B. B., & Decety, J. (2016). Callous traits in children with and without conduct problems predict reduced connectivity when viewing harm to others. *Scientific Reports*, 6(February), 2016.
- Zhang, J., Cao, W., Wang, M., Wang, N., Yao, S., & Huang, B. (2018). Multivoxel pattern analysis of structural MRI in children and adolescents with conduct disorder. *Brain Imaging and Behavior*, (Rubia 2011).
- Zhou, J., Yao, N., Fairchild, G., Cao, X., Zhang, Y., Xiang, Y. T., ... Wang, X. (2016). Disrupted default mode network connectivity in male adolescents with conduct disorder. *Brain Imaging and Behavior*.
- Zhou, J., Yao, N., Fairchild, G., Zhang, Y., & Wang, X. (2015). Altered hemodynamic activity in conduct disorder: A resting-state fMRI investigation. *PLoS ONE*, 10(3), 1–12.
- Zuckerman, M., & Neeb, M. (1979). Sensation seeking and psychopathology. *Psychiatry Research*, 1(3), 255–264.

Acknowledgments

First of all, I would like to thank my supervisor Prof. Dr. Daniel Brandeis. He gave me the opportunity to participate in the EU-projects Aggessotype and MATRICS and trusted me with my PhD in the framework of these ambitious and innovative studies. In particular, I would like to express my deepest gratitude for sharing his expertise, supporting the coordination of the study continuously, and promoting my academic development and successful completion of the present PhD thesis. Further, I would like to thank him for his enduring understanding for my concurrently duties and investments in the master of advanced studies in cognitive behavioral therapy and behavioral medicine at the University of Zurich.

Furthermore, many thanks go to all other members of our Aggessotype and MATRICS projects. I would like to express my sincere thanks to Prof. Dr. med. Dipl. Psych. Susanne Walitza for her tireless and extensive support, warm-hearted and motivating encouragement, and for having constant confidence in a successful course of our projects and my imbedded dissertation. Prof. Dr. Silvia Brem's constant interest and support was an elusively source of motivation. Above all, her helpful inputs further promote progress and eventually successful completion of the study and subsequent analyses. Especially at my start as a PhD student and study coordinator and during the first phases of the study, Dr. Juliane Ball accompanied and supported with endless effort whenever necessary. Further thanks belong to Dr. med. Anna Werling for their support regarding recruitment and PD Dr. Renate Drechsler for her help in implementing neuropsychological testing in our study. Dr. Jill Naaijen deserves immense thanks for being my unofficial mentor, answering all my questions with endless patience, and never forgetting to add some fun and a sense of humor regardless of the challenging situations now and then. In particular, I would like to thank Dr. Nathalie Holz for her continuous efforts and constructive comments and

inputs especially for my papers. Moreover, I am deeply thankful to M. Sc. Pascal Aggensteiner and his tireless engagement and reassuring support throughout the whole process of study coordination and academic process, not to mention the shared love for electronic music. My thanks also belong to Dr. Sarah Baumeister and Dr. Gabriela Gan for their scientific support and for always having an open ear. I am indebted to all the members of our Aggessotype and MATRICS projects for their important contributions to our project and for their important and constructive feedback on my papers.

Moreover, I am deeply grateful to Dr. Lukasz Smigielski, whose effortless scientific support and knowledge particularly in MR data analysis had an extensive effect on successfully accomplishing this PhD.

I would like to thank Dr. Philipp Stämpfli for his continuous support in the MR-Centre, particularly in implementing the MR sequences thoroughly and helping with his immense expertise in neuroimaging whenever we were confronted with any kind of problems during MR scanning, and Dr. Stefan Sommer for helping whenever necessary during the MR measurements.

I am thankful to Prof. Dr. Edna Grünblatt and her team for processing and storing our genetic samples, and eventually helped to transfer them to The Netherlands.

My thanks also belong to my colleagues M. Sc. Yu Jin Ressel, Dr. Alexander Roth, and our interns Noemi Signer and Aline Pfister for assisting in this project. Further, I would like to thank M. Sc. David Willinger and Dr. Iliana Karipidis for sharing their scientific knowledge and for their immense support and assistance when it came to practical issues with respect to MR analysis. My special thanks also go to M. Sc. Susanna Mohl. Without her tremendous assistance, this PhD in the framework of the Aggessotype and MATRICS projects would not have been possible. Moreover, I would like to thank M. Sc. Marco Hartmann, Dr. Georgette Pleisch, Dr. Isa Häberling, M. Sc. Sarah Di Pietro, M. Sc. Seline Coray, M. Sc. Sarah Di Pietro, M. Sc. Julia Strauss, M. Sc. Selina Beltrani, Dr. Gorka Fraga, and Dr. Franziska Minder for the pleasant company and enriching discussions during lunch and coffee breaks. I also want to thank Dr. Agnieszka Zuberer for the lively and reassuring conversations and her warm-hearted support not only in scientific but also private matters. Further, my thanks belong to M.Sc. Etna

Engeli, who taught me the first steps in conducting MR measurements and inspired me with her dedicated way of working.

Most importantly, my truly heart-felt thanks belong to my family and friends. Each of you made this PhD possible by supporting me in various ways, all characterized by endless support and understanding. The enriching, positive, and inspirational interactions with all of you extensively contributed to the successful completion of this dissertation.

Last but not least, to music and nature.

Curriculum Vitae

PERSONAL INFORMATION

Name: Julia Elisabeth Werhahn
Address: Schlossgasse 28
8003 Zürich
Date of birth: March 8th 1989
E-Mail: julia.werhahn@gmx.ch
Phone: +41 76 325 47 70
Nationality: German

EDUCATION

Since 09/2016	Doctoral studies in Psychology, University of Zurich
12/2015 – 05/2016	Qualified User Training, Magnetic Resonance Imaging Centre of the University Hospital of Psychiatry Zurich
Since 09/2014	Master of Advanced Studies in cognitive behavioral therapy and behavioral medicine, University of Zurich
10/2012 – 08/2014	Master of Science in Clinical Psychology, Neuro- and Rehabilitation Science, University of Freiburg (D)
09/2009 – 07/2012	Bachelor of Science in Psychology, University of Innsbruck (A)
09/2008 – 01/2012	Bachelor Study in Sociology, University of Innsbruck (A)

EMPLOYMENT HISTORY

01/2017 – 12/2018

Psychotherapist (in training), SNF-funded study of cognitive behavioral therapy for the generalized anxiety disorder, Department of Psychology, University of Zurich

Since 03/2016

Psychotherapist (in training), Ambulatory for cognitive behavioral therapy and behavioral medicine, Department of Psychology, University of Zurich

Since 02/2016

PhD Candidate Position, Department of Child and Adolescent Psychiatry and Psychotherapy, University Hospital of Psychiatry, University of Zurich

10/2014 – 01/2016

Postgraduate Psychologist, Department of Child and Adolescent Psychiatry and Psychotherapy, Pediatric Ward, Meilen

10/2013 – 04/2014

Student Assistant, Chair for Biological and Differential Psychology, University of Freiburg (D)

08 – 10/2013

Internship, Institute for Forensic Psychiatry and Psychotherapy, Langenthal

05 – 07/2013

Voluntary Scientific Internship, Research- and Innovation Project for the Development of a Multimodal Online Stress Prevention Training, Chair for Biological and Differential Psychology, University of Freiburg (D)

11/2012 – 07/2013

Student Assistant, Department for Quality Management and Social Medicine, University Hospital Freiburg (D)

- 04 – 06/2012 **Voluntary Scientific Internship**, Chair for Social Psychology, Institute of Psychology, University of Innsbruck (A)
- 09 – 10/2011 **Staff**, 10th International Disaster Management Meeting, Austrian Red Cross, Innsbruck (A)
- 07 – 09/2011 **Internship**, Department of Pediatrics and Adolescent Medicine, University Hospitals Innsbruck (A)
- 02/2011 **Internship**, Social-Pediatric Centre, Children`s Hospital Garmisch-Partenkirchen (D)

CONFERENCES AND EU-PROJECT MEETINGS

- 10 – 11/2016 **General Assembly Meeting** of the EU-funded project “Aggressotype”, Laptop Presentation, Nijmegen & Utrecht (NL)
- 09/2017 **13th Symposium of the Zurich Center for Integrative Human Physiology (ZIHP)**, Poster Presentation
- 09/2017 **Neuroscience Center Zurich (ZNZ) Symposium 2017**, Poster Presentation
- 11/2017 **4th Burghölzli Meeting**, Poster Presentation, University Hospital of Psychiatry, Zurich
- 11/2017 **General Assembly Meeting** of the EU-funded project “Aggressotype”, Poster and Laptop Presentation, Fuerteventura (ES)
- 08/2018 **14th Symposium of the Zurich Center for Integrative Human Physiology (ZIHP)**, Poster Presentation

-
- 09/2018 **General Assembly Meeting** of the EU-funded project “Aggressotype”, Poster Presentation, Rome (IT)
- 09/2018 **Neuroscience Center Zurich (ZNZ) Symposium 2018**, Poster Presentation
- 10/2018 **5th Burghölzli Meeting**, Oral Contribution and Poster Presentation, University Hospital of Psychiatry, Zurich